

GENERALIZED QUASILINEARIZATION METHOD FOR A  
FORCED DUFFING EQUATION WITH MIXED NONLINEAR  
THREE-POINT BOUNDARY CONDITIONS

Ahmed Alsaedi

Department of Mathematics

Faculty of Science

King Abdul Aziz University

P.O. Box. 80257, Jeddah, 21589, KINGDOM OF SAUDI ARABIA

e-mail: aalsaedi@hotmail.com

**Abstract:** In this paper, we apply the generalized quasilinearization technique to a three-point nonlinear mixed boundary value problem involving a forced Duffing equation and obtain sequences of upper and lower solutions converging monotonically and quadratically to the unique solution of the problem.

**AMS Subject Classification:** 34B10, 34B15

**Key Words:** quasilinearization, Duffing equation, three-point boundary value problem, quadratic convergence

## 1. Introduction

Duffing equation is a well known nonlinear equation of applied science which is used as a powerful tool to discuss some important practical phenomena such as periodic orbit extraction, nonuniformity caused by an infinite domain, nonlinear mechanical oscillators, etc. Another important application of Duffing equation is in the field of the prediction of diseases. A careful measurement and analysis of a strongly chaotic voice has the potential to serve as an early warning system for more serious chaos and possible onset of disease. This chaos is stimulated

with the help of Duffing equation. In fact, the success at analyzing and predicting the onset of chaos in speech and its simulation by equations such as the Duffing equation has enhanced the hope that we might be able to predict the onset of arrhythmia and heart attacks someday. Such predictions are based on the numerical solutions of the Duffing equation. However, there do exist a number of powerful procedures for obtaining approximate solutions of nonlinear problems such as Newton-Raphson method, Galerkins method, expansion methods, iterative techniques, method of upper and lower solutions to name a few. The monotone iterative method and Newton's method are known to be two efficient techniques for finding roots of nonlinear equations. The first one applies to equations involving monotone operators and produces a sequence converging monotonically to a solution. The Newton's method has the advantage over the monotone iterative method that it provides quadratically convergent sequences. Applied to nonlinear differential equations, Newton's method is known as the quasilinearization method. The origin of the quasilinearization lies in the theory of dynamic programming Bellman [6], Bellman and Kalaba [7] and Lee [20]. This method applies to semilinear equations with convex or concave nonlinearities and provides an explicit analytic representation of approximate solution of the given problem. However, the concavity/convexity assumption proved to be a stumbling block for further development of the theory. The nineties brought new dimensions to this technique. The most interesting new idea was introduced by Lakshmikantham [17-18] who generalized the method of quasilinearization by relaxing the convexity assumption. This extension, now known as generalized quasilinearization, consists of the method of lower and upper solutions and monotone iterative technique together with differential inequalities and comparison results. This development was so significant that it attracted the attention of many researchers and the method was extensively developed and applied to a wide range of initial and boundary value problems Ahmad [1], Ahmad et al [3-4], Buica [8], Cabada and Nieto [9], Lakshmikantham and Vatsala [19] and references therein. Some real-world applications of the quasilinearization technique can be found in Mandelzweig and Tabakin [23], Nieto and Torres [24], Nikolov et al [25], and Yermachenko and Sadyrbaev [26].

Multi-point nonlinear boundary value problems, which refer to a different family of boundary conditions in the study of disconjugacy theory Coppel [10], have been addressed by many authors, for example, Kiguradze and Lomtadze [16], Gupta [13], Gupta and Trofimchuck [14], Ma [20-21], Bai and Fang [5], and Eloe and Ahmad [11]. Eloe and Gao [12] discussed the quasilinearization method for a three-point boundary value problem. Ahmad [2] developed the generalized quasilinearization method for a general three-point nonlinear

boundary value problem.

In this paper, we consider a forced Duffing equation with nonlinear three-point mixed boundary conditions and develop a monotone iteration scheme by relaxing the convexity assumption on the function involved in the differential equation and the concavity assumption on nonlinearities in the boundary conditions. In fact, we obtain monotone sequences of iterates (approximate solutions) converging quadratically to the unique solution of the three-point boundary value problem.

### 2. Preliminaries

We consider a three-point boundary value problem for the forced Duffing equation given by

$$x'' + kx' + f(t, x) = 0, \tag{2.1}$$

$$px(0) - qx'(0) = g_1(x(1/2)), \quad px(1) + qx'(1) = g_2(x(1/2)), \tag{2.2}$$

where  $f$  is continuous with  $f_x < 0$  on  $[0, 1] \times R$ ,  $k \in R$  such that  $k \neq 0$ ,  $p, q > 0$  with  $p > 1$  and  $g_i : R \rightarrow R$ ,  $i = 1, 2$  are continuous.

By Green's function method, the solution,  $x(t)$  of (2.1)-(2.2) can be written as

$$\begin{aligned} x(t) &= g_1(x(1/2)) \left[ \frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]} \right] \\ &+ g_2(x(1/2)) \left[ \frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]} \right] + \int_0^1 G_k(t, s) f(s, x(s)) ds, \end{aligned}$$

where

$$G_k(t, s) = \begin{cases} \sigma \left[ \frac{p - kq}{p} - e^{k(1-s)} \right] \left[ \frac{p + kq}{p} - e^{-kt} \right], & 0 \leq t \leq s, \\ \sigma \left[ \frac{p - kq}{p} - e^{k(1-t)} \right] \left[ \frac{p + kq}{p} - e^{-ks} \right], & s \leq t \leq 1, \end{cases}$$

$$\sigma = \frac{pe^{ks}}{k[(p - kq) - (p + kq)e^k]}.$$

We say that  $\alpha \in C^2[0, 1]$  is a lower solution of the boundary value problem (2.1)-(2.2) if

$$\begin{aligned} \alpha''(t) + k\alpha'(t) + f(t, \alpha(t)) &\geq 0, \quad t \in [0, 1], \\ p\alpha(0) - q\alpha'(0) &\leq g_1(\alpha(1/2)), \quad p\alpha(1) + q\alpha'(1) \leq g_2(\alpha(1/2)), \end{aligned}$$

and  $\beta \in C^2[0, 1]$  is an upper solution of (2.1)-(2.2) if

$$\begin{aligned} \beta''(t) + k\beta'(t) + f(t, \beta(t)) &\leq 0, \quad t \in [0, 1], \\ p\beta(0) - q\beta'(0) &\geq g_1(\beta(1/2)), \quad p\beta(1) + q\beta'(1) \geq g_2(\beta(1/2)). \end{aligned}$$

Now, we present comparison and existence results related to (2.1)-(2.2) which play a pivotal role in proving the main result.

**Theorem 2.1.** *Assume that  $f$  is continuous with  $f_x < 0$  on  $[0, 1] \times R$  and  $g$  is continuous on  $R$  satisfying a one-sided Lipschitz condition:*

$$g_i(x) - g_i(y) \leq L_i(x - y), \quad 0 \leq L_i < 1, \quad i = 1, 2.$$

*Let  $\beta$  and  $\alpha$  be the upper and lower solutions of (2.1)-(2.2) respectively. Then  $\alpha(t) \leq \beta(t)$ ,  $t \in [0, 1]$ .*

*Proof.* Define  $h(t) = \alpha(t) - \beta(t)$ . For the sake of contradiction, we suppose that  $h(t) > 0$  for some  $t \in [0, 1]$ . First we take  $t_0 \in (0, 1)$ . Then by the definition of lower and upper solutions and the assumption  $f_x < 0$ , we obtain

$$\begin{aligned} h''(t_0) + kh'(t_0) &= \alpha''(t_0) + k\alpha'(t_0) - \beta''(t_0) - k\beta'(t_0) \\ &\geq -f(t_0, \alpha(t_0)) + f(t_0, \beta(t_0)) > 0. \end{aligned}$$

Now, employing a standard procedure Jackson [15] in the applications of upper and lower solutions, let  $h(t)$  have a local positive maximum at  $t_0 \in (0, 1)$ , then  $h'(t_0) = 0$  and  $h''(t_0) \leq 0$ , which contradicts the above inequality. Thus, for  $t_0 \in (0, 1)$ , we have  $\alpha(t) \leq \beta(t)$ . Now, suppose that  $h(t)$  has a local positive maximum at  $t_0 = 1$ , then  $h'(1) = 0$  and  $h''(1) < 0$ . On the other hand, using the definition of lower and upper solutions together with the fact that  $g_2$  satisfies a one sided Lipschitz condition, we find that

$$\begin{aligned} ph(1) + qh'(1) &= p\alpha(1) + q\alpha'(1) - (p\beta(1) + q\beta'(1)) \leq g_2(\alpha(\frac{1}{2})) - g_2(\beta(\frac{1}{2})) \\ &< \alpha(\frac{1}{2}) - \beta(\frac{1}{2}) = h(\frac{1}{2}). \end{aligned}$$

Thus  $ph(1) < h(\frac{1}{2})$  or  $h(1) < h(\frac{1}{2})$  for  $p > 1$ , which is a contradiction. Similarly, we get a contradiction for  $t_0 = 0$ . Hence we conclude that  $\alpha(t) \leq \beta(t)$  on  $[0, 1]$ . □

**Theorem 2.2.** *Assume that  $f$  is continuous on  $[0, 1] \times R$  with  $f_x < 0$  and  $g_i$  are continuous on  $R$  satisfying one-sided Lipschitz condition:*

$$g_i(x) - g_i(y) \leq L_i(x - y), \quad 0 \leq L_i < 1, \quad i = 1, 2.$$

*Further, we assume that there exist an upper solution  $\beta$  and a lower solution  $\alpha$  of (2.1)-(2.2) such that  $\alpha(t) \leq \beta(t)$ ,  $t \in [0, 1]$ . Then there exists a solution  $x(t)$  of (2.1)-(2.2) satisfying  $\alpha(t) \leq x(t) \leq \beta(t)$ ,  $t \in [0, 1]$ .*

*Proof.* Let us define  $F$  and  $G$  by

$$F(t, x) = \begin{cases} f(t, \beta) - \frac{x-\beta}{1+x-\beta}, & \text{if } x(t) > \beta(t), \\ f(t, x), & \text{if } \alpha(t) \leq x(t) \leq \beta(t), \\ f(t, \alpha) - \frac{x-\alpha}{1+|x-\alpha|}, & \text{if } x(t) < \alpha(t), \end{cases}$$

$$\hat{g}_i(x) = \begin{cases} g_i(\beta(\frac{1}{2})), & \text{if } x > \beta(\frac{1}{2}), \\ g_i(x), & \text{if } \alpha(\frac{1}{2}) \leq x \leq \beta(\frac{1}{2}), \\ g_i(\alpha(\frac{1}{2})), & \text{if } x < \alpha(\frac{1}{2}), \end{cases}$$

for  $i = 1, 2$ .

Since  $F(t, x)$  and  $\hat{g}_i(x)$  are continuous and bounded, a standard application of Schauder’s Fixed Point Theorem ensures the existence of a solution,  $x$  of the problem

$$x''(t) + kx'(t) + F(t, x(t)) = 0, \quad t \in [0, 1],$$

$$px(0) - qx'(0) = \hat{g}_1(x(1/2)), \quad px(1) + qx'(1) = \hat{g}_2(x(1/2)).$$

In order to complete the proof, we need to show that  $\alpha(t) \leq x(t) \leq \beta(t)$  on  $[0, 1]$ . For that, we set  $h(t) = \alpha(t) - x(t)$ . For the sake of the contradiction, let  $h(t) > 0$  for some  $t \in [0, 1]$ . We define

$$t_0 = \inf\{\tau \in [0, 1] : h(\tau) \geq h(t), 0 \leq t \leq 1\},$$

and note that  $0 < t_0$  by continuity. As  $\hat{g}_2$  satisfies a one-sided Lipschitz condition on  $[\alpha(\frac{1}{2}), \beta(\frac{1}{2})]$ , it follows that

$$\begin{aligned} ph(1) + qh'(1) &= p\alpha(1) + q\alpha'(1) - (px(1) + qx'(1)) \leq \hat{g}_2(\alpha(1/2)) - \hat{g}_2(x(1/2)) \\ &< (\alpha(1/2) - x(1/2)) = h(1/2). \end{aligned}$$

As in the proof of Theorem 2.1, let  $h(t)$  have a local maximum at  $t_0 \in (0, 1)$  implying that  $h'(t_0) = 0$  and  $h''(t_0) \leq 0$ . On the other hand, by the definition of upper and lower solutions together with the assumption  $F_x < 0$ , we have

$$\begin{aligned} h''(t_0) + kh'(t_0) &= \alpha''(t_0) + k\alpha'(t_0) - (x''(t_0) + kx'(t_0)) \\ &\geq -F(t_0, \alpha(t_0)) + F(t_0, x(t_0)) > 0. \end{aligned}$$

This contradicts our supposition. Hence  $\alpha(t) - x(t) \leq 0$ . Similarly, it can be shown that  $x(t) \leq \beta(t)$ . Thus, it follows that  $\alpha(t) \leq x(t) \leq \beta(t)$ ,  $t \in [0, 1]$ .  $\square$

### 3. Main Result

**Theorem 3.** *Assume that:*

(A<sub>1</sub>)  $\alpha_0, \beta_0$  are lower and upper solutions of (2.1)-(2.2) respectively.

(A<sub>2</sub>)  $f(t, x) \in C([0, 1] \times R)$  be such that  $f_x < 0$  and  $(f_{xx}(t, x) + \phi_{xx}(t, x)) \geq 0$ , where  $\phi_{xx}(t, x) \geq 0$  for some continuous function  $\phi(t, x)$  on  $[0, 1] \times R$ .

(A<sub>3</sub>) For  $i = 1, 2$ ,  $g_i(x), g'_i(x), g''_i(x)$  are continuous on  $R$  with  $0 \leq g'_i \leq 1$  and  $g''_i(x) + \psi''_i(x) \leq 0$  with  $\psi''_i \leq 0$  on  $R$  for some continuous functions  $\psi_i(x)$ .

Then there exist monotone sequences  $\{\alpha_n\}, \{\beta_n\}$  that converge quadratically in the space of continuous functions on  $[0, 1]$  to the unique solution  $x$  of (2.1)-(2.2).

*Proof.* Define  $F : [0, 1] \times R \rightarrow R$  by

$$F(t, x) = f(t, x) + \phi(t, x),$$

and  $G_i : R \rightarrow R$  by

$$G_i(x) = g_i(x) + \psi_i(x), \quad i = 1, 2.$$

Using the Generalized Mean Value Theorem together with (A<sub>2</sub>) and (A<sub>3</sub>), we obtain

$$f(t, x) \geq f(t, y) + F_x(t, y)(x - y) + \phi(t, y) - \phi(t, x), \tag{3.1}$$

$$g_i(x) \leq g_i(y) + G'_i(y)(x - y) + \psi_i(y) - \psi_i(x), \quad i = 1, 2. \tag{3.2}$$

Now, we set

$$F(t, x; \alpha_0) = f(t, \alpha_0) + F_x(t, \alpha_0)(x - \alpha_0) + \phi(t, \alpha_0) - \phi(t, x),$$

$$\overline{F}(t, x; \alpha_0, \beta_0) = f(t, \beta_0) + F_x(t, \alpha_0)(x - \beta_0) + \phi(t, \beta_0) - \phi(t, x),$$

and

$$\begin{aligned} h_i(x(1/2); \alpha_0, \beta_0) &= g_i(\alpha_0(1/2)) + G'_i(\beta_0(1/2))(x(1/2) - \alpha_0(1/2)) \\ &\quad + \psi_i(\alpha_0(1/2)) - \psi_i(x(1/2)), \end{aligned}$$

$$\begin{aligned} \hat{h}_i(x(1/2); \beta_0) &= g_i(\beta_0(1/2)) + G'_i(\beta_0(1/2))(x(1/2) - \beta_0(1/2)) \\ &\quad + \psi_i(\beta_0(1/2)) - \psi_i(x(1/2)), \quad i = 1, 2. \end{aligned}$$

We now consider the BVPs

$$x''(t) + kx'(t) + F(t, x; \alpha_0) = 0, \quad t \in [0, 1], \tag{3.3}$$

$$px(0) - qx'(0) = h_1(x(1/2); \alpha_0, \beta_0), \quad px(1) + qx'(1) = h_2(x(1/2); \alpha_0, \beta_0), \tag{3.4}$$

and

$$x''(t) + kx'(t) + \overline{F}(t, x; \alpha_0, \beta_0) = 0, \quad t \in [0, 1], \tag{3.5}$$

$$px(0) - qx'(0) = \hat{h}_1(x(1/2), \beta_0), \quad px(1) + qx'(1) = \hat{h}_2(x(1/2), \beta_0). \tag{3.6}$$

Let us show that  $\alpha_0$  and  $\beta_0$  are respectively lower and upper solutions of (3.3)-(3.4). By definition of lower solution and the fact that  $F(t, \alpha_0; \alpha_0) = f(t, \alpha_0)$ ,

we get

$$\begin{aligned} \alpha_0'' + k\alpha_0' + F(t, \alpha_0; \alpha_0) &= \alpha_0'' + k\alpha_0' + f(t, \alpha_0) \geq 0, \\ p\alpha_0(0) - q\alpha_0'(0) &\leq g_1(\alpha_0(1/2)) = h_1(\alpha_0(1/2); \alpha_0; \beta_0), \\ p\alpha_0(1) + q\alpha_0'(1) &\leq g_2(\alpha_0(1/2)) = h_2(\alpha_0(1/2); \alpha_0; \beta_0), \end{aligned}$$

which implies that  $\alpha_0$  is a lower solution of (3.3)-(3.4). Using (3.1) and the definition of upper solution, we have

$$\begin{aligned} &\beta_0'' + k\beta_0' + F(t, \beta_0; \alpha_0) \\ &= \beta_0'' + k\beta_0' + f(t, \alpha_0) + F_x(t, \alpha_0)(\beta_0 - \alpha_0) + \phi(t, \alpha_0) - \phi(t, \beta_0) \\ &\leq \beta_0'' + k\beta_0' + f(t, \beta_0) \leq 0. \end{aligned}$$

Using mean value theorem and the nonincreasing property of  $G_1'$ , we have

$$\begin{aligned} &g_1(\beta_0(1/2)) - h_1(\beta_0(1/2); \alpha_0, \beta_0) \\ &= g_1(\beta_0(1/2)) - g_1(\alpha_0(1/2)) - G_1'(\beta_0(1/2))(\beta_0(1/2) - \alpha_0(1/2)) \\ &\quad - \psi_1(\alpha_0(1/2)) + \psi_1(\beta_0(1/2)) \\ &= G_1(\beta_0(1/2)) - G_1(\alpha_0(1/2)) - G_1'(\beta_0(1/2))(\beta_0(1/2) - \alpha_0(1/2)) \\ &= [G_1'(c_0) - G_1'(\beta_0(1/2))](\beta_0(1/2) - \alpha_0(1/2)) \geq 0, \end{aligned}$$

where  $\alpha_0(1/2) \leq c_0 \leq \beta_0(1/2)$ . Consequently, we have

$$p\beta_0(0) - q\beta_0'(0) \geq h_1(\beta_0(1/2); \alpha_0, \beta_0).$$

Similarly, it can be shown that

$$p\beta_0(1) + q\beta_0'(1) \geq h_2(\beta_0(1/2); \alpha_0, \beta_0).$$

Thus,  $\beta_0$  is an upper solution of (3.3)-(3.4). Hence, by Theorem 2.2, there is a solution  $\alpha_1$  of (3.3)-(3.4) satisfying

$$\alpha_0(t) \leq \alpha_1(t) \leq \beta_0(t), \quad t \in [0, 1]. \tag{3.7}$$

Note that Theorem 2.2 applies since  $h_i' = g_i'(\beta_0(1/2))$ ,  $i = 1, 2$ . Similarly,  $\beta_0$  is an upper solution of (3.5)-(3.6) as

$$\begin{aligned} \overline{F}(t, \beta_0; \alpha_0; \beta_0) &= f(t, \beta_0), \quad g_1(\beta_0(1/2)) = \hat{h}_1(\beta_0(1/2); \beta_0), \\ g_2(\beta_0(1/2)) &= \hat{h}_2(\beta_0(1/2); \beta_0). \end{aligned}$$

As before, using (3.1), we obtain

$$\begin{aligned} \alpha_0'' + k\alpha_0' + \overline{F}(t, \alpha_0; \alpha_0, \beta_0) &= \alpha_0'' + k\alpha_0' + f(t, \beta_0) + F_x(t, \alpha_0)(\alpha_0 - \beta_0) \\ &\quad + \phi(t, \beta_0) - \phi(t, \alpha_0) \geq \alpha_0'' + k\alpha_0' + f(t, \alpha_0) \geq 0. \end{aligned}$$

Now, we will show that  $p\alpha_0(0) - q\alpha_0'(0) \leq \hat{h}_1(\alpha_0(1/2); \beta_0)$ . By Mean Value Theorem, we find that

$$\hat{h}_1(\alpha_0(1/2); \beta_0) - g_1(\alpha_0(1/2)) = g_1(\beta_0(1/2)) + G_1'(\beta_0(1/2))(\alpha_0(1/2) - \beta_0(1/2))$$

$$\begin{aligned}
 & + \psi_1(\beta_0(1/2)) - \psi_1(\alpha_0(1/2)) - g_1(\alpha_0(1/2)) \\
 & = G_1(\beta_0(1/2)) - G_1(\alpha_0(1/2)) + G'_1(\beta_0(1/2))(\alpha_0(1/2) - \beta_0(1/2)) \\
 & = [G'_1(c_1) - G'_1(\beta_0(1/2))](\beta_0(1/2) - \alpha_0(1/2)) \geq 0,
 \end{aligned}$$

where  $\alpha_0(1/2) \leq c_1 \leq \beta_0(1/2)$ . Thus

$$p\alpha_0(0) - q\alpha'_0(0) \leq g_1(\alpha_0(1/2)) \leq \hat{h}_1(\alpha_0(1/2); \beta_0).$$

Similarly, it can be shown that

$$p\alpha_0(1) + q\alpha'_0(1) \leq \hat{h}_2(\alpha_0(1/2); \beta_0).$$

Thus,  $\alpha_0$  is a lower solution of (3.5)-(3.6). Again, by Theorem 2.2, there exists a solution  $\beta_1$  of (3.5)-(3.6) such that

$$\alpha_0(t) \leq \beta_1(t) \leq \beta_0(t), \quad t \in [0, 1]. \tag{3.8}$$

Now, we show that  $\alpha_1 \leq \beta_1$ . To do this we prove that  $\alpha_1, \beta_1$  are lower and upper solutions of (2.1)-(2.2) respectively. Using the fact that  $\alpha_1$  is a solution of (3.3)-(3.4), we get

$$\begin{aligned}
 & \alpha''_1(t) + k\alpha'_1(t) + f(t, \alpha_1) \\
 & \geq \alpha''_1(t) + k\alpha'_1(t) + f(t, \alpha_0) + F_x(t, \alpha_0)(\alpha_1 - \alpha_0) + \phi(t, \alpha_0) - \phi(t, \alpha_1) \\
 & = \alpha''_1(t) + k\alpha'_1(t) + F(t, \alpha_1; \alpha_0) = 0.
 \end{aligned}$$

Now, in view of nonincreasing property of  $G'_1$ , we obtain

$$\begin{aligned}
 & g_1(\alpha_1(1/2)) - [p\alpha_1(0) - q\alpha'_1(0)] \\
 & = g_1(\alpha_1(1/2)) - g_1(\alpha_0(1/2)) - G'_1(\beta_0(1/2))(\alpha_1(1/2) - \alpha_0(1/2)) \\
 & - \psi_1(\alpha_0(1/2)) + \psi_1(\alpha_1(1/2)) = [G'_1(c_2) - G'_1(\beta_0(1/2))](\alpha_1(1/2) - \alpha_0(1/2)) \geq 0,
 \end{aligned}$$

where  $c_2 \in (\alpha_0(1/2), \alpha_1(1/2))$ , which in turn yields

$$p\alpha_1(0) - q\alpha'_1(0) \leq g_1(\alpha_1(1/2)).$$

Similarly, it can be shown that  $p\alpha_1(1) + q\alpha'_1(1) \leq g_2(\alpha_1(1/2))$ . This implies that  $\alpha_1$  is a lower solution of (2.1)-(2.2). Similarly, it can be shown that  $\beta_1$  is an upper solution of (2.1)-(2.2). By Theorem 2.1, it follows that

$$\alpha_1(t) \leq \beta_1(t), \quad t \in [0, 1]. \tag{3.9}$$

Combining (3.7), (3.8) and (3.9) yields

$$\alpha_0(t) \leq \alpha_1(t) \leq \beta_1(t) \leq \beta_0(t), \quad t \in [0, 1].$$

Continuing this process, by induction, one can prove that

$$\alpha_n(t) \leq \alpha_{n+1}(t) \leq \beta_{n+1}(t) \leq \beta_n(t), \quad t \in [0, 1], \quad n = 0, 1, \dots,$$

where  $\alpha_{n+1}$  satisfies the problem

$$x''(t) + kx'(t) + F(t, x; \alpha_n) = 0, \quad t \in [0, 1],$$



$$px(0) - qx'(0) = h_1(x(1/2); \alpha_n, \beta_n), \quad px(1) + qx'(1) = h_2(x(1/2); \alpha_n, \beta_n)$$

and  $\beta_{n+1}$  satisfies the BVP

$$x''(t) + kx'(t) + \bar{F}(t, x; \alpha_n, \beta_n) = 0, \quad t \in [0, 1],$$

$$px(0) - qx'(0) = \hat{h}_1(x(1/2); \beta_n), \quad px(1) + qx'(1) = \hat{h}_2(x(1/2); \beta_n).$$

Since  $[0, 1]$  is compact and the convergence is monotone, it follows that the convergence of each sequence  $\{\alpha_n\}$  and  $\{\beta_n\}$  is uniform. Employing the standard arguments Lakshmikantham and Vatsala [19], we conclude that  $x$  is the limit point of each of the two sequences and consequently, we get

$$\begin{aligned} x(t) &= g_1(x(1/2)) \left[ \frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]} \right] \\ &+ g_2(x(1/2)) \left[ \frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]} \right] + \int_0^1 G_k(t, s) f(s, x(s)) ds. \end{aligned}$$

This proves that  $x$  is the unique solution of (2.1)-(2.2).

In order to prove that each of the sequences  $\{\alpha_n\}, \{\beta_n\}$  converges quadratically, we set  $q_n = \beta_n - x \geq 0, \quad p_n = x - \alpha_n \geq 0$ , where  $x$  denotes the unique solution of (2.1)-(2.2). We only show the quadratic convergence with  $p_n$  as the details for the quadratic convergence for  $q_n$  are similar. Applying the mean value theorem, there exist  $\alpha_n \leq c_3, c_4, c_5 \leq x$  and  $\alpha_n \leq \zeta_1 \leq \alpha_{n+1}$  such that

$$\begin{aligned} &p''_{n+1} + kp'_{n+1} \\ &= -f(t, x) + f(t, \alpha_n) + F_x(t, \alpha_n)(\alpha_{n+1} - \alpha_n) + \phi(t, \alpha_n) - \phi(t, \alpha_{n+1}) \\ &= -f_x(t, c_3)(x - \alpha_n) + F_x(t, \alpha_n)(\alpha_{n+1} - x + x - \alpha_n) - \phi_x(t, \zeta_1)(\alpha_{n+1} - \alpha_n) \\ &= [-F_x(t, c_3) + F_x(t, \alpha_n) + \phi_x(t, c_3) - \phi_x(t, \zeta_1)]p_n + [-F_x(t, \alpha_n) + \phi_x(t, \zeta_1)]p_{n+1} \\ &\geq [-F_x(t, x) + F_x(t, \alpha_n) + \phi_x(t, \alpha_n) - \phi_x(t, x)]p_n + [-F_x(t, \zeta_1) + \phi_x(t, \zeta_1)]p_{n+1} \\ &= -F_{xx}(t, c_4)p_n^2 - \phi_{xx}(t, c_5)p_n^2 - f_x(t, \zeta_1)p_{n+1} \\ &\geq -M\|p_n\|^2, \end{aligned}$$

where  $A$  is a bound on  $\|F_{xx}\|, B$  is a bound on  $\|\phi_{xx}\|$  for  $t \in [0, 1]$  and  $M = A + B$ . Here  $\|\cdot\|$  denotes the supremum norm on  $C[0, 1]$ . Also there exist  $\alpha_n(1/2) \leq c_6, r_1 \leq c_7, r_2 \leq x \leq \beta_n$  and  $\alpha_n \leq \zeta_2, \zeta_3 \leq \zeta_4, \zeta_5 \leq \alpha_{n+1}$  such that

$$\begin{aligned} p_{n+1}(t) &= [g_1(x(1/2)) - h_1(\alpha_{n+1}(1/2); \alpha_n, \beta_n)] \left( \frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]} \right) \\ &+ [g_2(x(1/2)) - h_2(\alpha_{n+1}(1/2); \alpha_n, \beta_n)] \left( \frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]} \right) \\ &+ \int_0^1 G_k(t, s) [f(s, x) - F(s, \alpha_{n+1}; \alpha_n)] ds \end{aligned}$$

$$\begin{aligned}
&= [g_1(x(1/2)) - g_1(\alpha_n(1/2) - G'_1(\beta_n(1/2))(\alpha_{n+1}(1/2) - \alpha_n(1/2)) \\
&\quad - \psi_1(\alpha_n(1/2)) + \psi_1(\alpha_{n+1}(1/2))]\left(\frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]}\right) \\
&+ [g_2(x(1/2)) - g_2(\alpha_n(1/2) - G'_2(\beta_n(1/2))(\alpha_{n+1}(1/2) - \alpha_n(1/2)) \\
&\quad - \psi_2(\alpha_n(1/2)) + \psi_2(\alpha_{n+1}(1/2))]\left(\frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]}\right) \\
&\quad - \int_0^1 G_k(t, s)[p''_{n+1} + kp'_{n+1}]ds \\
&= [g'_1(c_6)(x(1/2) - \alpha_n(1/2)) - G'_1(\beta_n(1/2))(\alpha_{n+1} - \alpha_n(1/2)) \\
&\quad + \psi'_1(\zeta_2)(\alpha_{n+1}(1/2) - \alpha_n(1/2))]\left(\frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]}\right) \\
&+ [g'_2(r_1)(x(1/2) - \alpha_n(1/2)) - G'_2(\beta_n(1/2))(\alpha_{n+1} - \alpha_n(1/2)) \\
&\quad + \psi'_2(\zeta_3)(\alpha_{n+1}(1/2) - \alpha_n(1/2))]\left(\frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]}\right) \\
&\quad - \int_0^1 G_k(t, s)[p''_{n+1} + kp'_{n+1}]ds \\
&\leq [(G'_1(c_6) - G'_1(\beta_n(1/2)) - (\psi'_1(c_6) - \psi'_1(\zeta_2)))p_n(1/2) \\
&+ (G'_1(\beta_n(1/2)) - \psi'_1(\zeta_2))p_{n+1}(1/2)]\left(\frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]}\right) \\
&\quad + [(G'_2(r_1) - G'_2(\beta_n(1/2)) - (\psi'_2(r_1) - \psi'_2(\zeta_3)))p_n(1/2) \\
&+ (G'_2(\beta_n(1/2)) - \psi'_2(\zeta_3))p_{n+1}(1/2)]\left(\frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]}\right) \\
&\quad + M\|p_n\|^2 \int_0^1 |G_k(t, s)|ds \\
&\leq [-G''_1(c_7)(\beta_n(1/2) - c_6)p_n(1/2) - \psi''_1(\zeta_4)p_n^2(1/2) \\
&\quad + g'_1(\beta_n(1/2))p_{n+1}(1/2)]\left(\frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]}\right) \\
&\quad + [-G''_2(r_2)(\beta_n(1/2) - r_1)p_n(1/2) - \psi''_2(\zeta_5)p_n^2(1/2) \\
&+ g'_2(\beta_n(1/2))p_{n+1}(1/2)]\left(\frac{(p + kq) - pe^{-kt}}{p[(p + kq) - (p - kq)e^{-k}]}\right) + M_1\|p_n\|^2 \\
&\leq [-G''_1(c_7)(\beta_n(1/2) - \alpha_n(1/2))p_n(1/2) - \psi''_1(\zeta_4)p_n^2(1/2) \\
&\quad + g'_1(\beta_n(1/2))p_{n+1}(1/2)]\left(\frac{(p - kq)e^{-k} - pe^{-kt}}{p[(p - kq)e^{-k} - (p + kq)]}\right)
\end{aligned}$$

$$\begin{aligned}
 &+ [-G_2''(r_2)(\beta_n(1/2) - \alpha_n(1/2))p_n(1/2) - \psi_2''(\zeta_5)p_n^2(1/2) \\
 &+ g_2'(\beta_n(1/2))p_{n+1}(1/2)]\left(\frac{(p+kq) - pe^{-kt}}{p[(p+kq) - (p-kq)e^{-k}]}\right) + M_1\|p_n\|^2 \\
 &= [-G_1''(c_7)(q_n(1/2) + p_n(1/2))p_n(1/2) - \psi_1''(\zeta_4)p_n^2(1/2) \\
 &\quad + g_1'(\beta_n(1/2))p_{n+1}(1/2)]\left(\frac{(p-kq)e^{-k} - pe^{-kt}}{p[(p-kq)e^{-k} - (p+kq)]}\right) \\
 &+ [-G_2''(r_2)(q_n(1/2) + p_n(1/2))p_n(1/2) - \psi_2''(\zeta_5)p_n^2(1/2) \\
 &+ g_2'(\beta_n(1/2))p_{n+1}(1/2)]\left(\frac{(p+kq) - pe^{-kt}}{p[(p+kq) - (p-kq)e^{-k}]}\right) + M_1\|p_n\|^2 \\
 &\leq [N_1\left(\frac{1}{2}q_n^2(1/2) + \frac{3}{2}p_n^2(1/2)\right) + g_1'(\beta_n(1/2))p_{n+1}(1/2) + D_1p_n^2(1/2)]N_2 \\
 &+ [M_2\left(\frac{1}{2}q_n^2(1/2) + \frac{3}{2}p_n^2(1/2)\right) + g_2'(\beta_n(1/2))p_{n+1}(1/2) + D_2p_n^2(1/2)]M_3 \\
 &\quad + M_1\|p_n\|^2 \leq \left[\left(\frac{3}{2}N_1 + D_1\right)p_n^2(1/2) + \frac{N_1}{2}q_n^2(1/2) + \lambda_1p_{n+1}(1/2)\right]N_2 \\
 &\quad + \left[\left(\frac{3}{2}M_2 + D_2\right)p_n^2(1/2) + \frac{M_2}{2}q_n^2(1/2) + \lambda_2p_{n+1}(1/2)\right]M_3 + M_1\|p_n\|^2 \\
 &\leq \left(\frac{3}{2}N_1N_2 + D_1N_2 + \frac{3}{2}M_2M_3 + D_2M_3 + M_1\right)\|p_n\|^2 \\
 &\quad + \left(\frac{N_1}{2}N_2 + \frac{M_2}{2}M_3\right)\|q_n\|^2 + (\lambda_1 + \lambda_2)\|p_{n+1}\|,
 \end{aligned}$$

where  $|g_1'| \leq \lambda_1 < 1$ ,  $|g_2'| \leq \lambda_2 < 1$ ,  $|G_1''| < N_1$ ,  $|G_2''| < M_2$ ,  $|\psi_1''| < D_1$ ,  $|\psi_2''| < D_2$ ,  $\left|\frac{(p-kq)e^{-k} - pe^{-kt}}{p[(p-kq)e^{-k} - (p+kq)]}\right| < N_1$ ,  $\left|\frac{(p+kq) - pe^{-kt}}{p[(p+kq) - (p-kq)e^{-k}]}\right| < M_3$  and  $M_1$  provides a bound on  $M \int_0^1 |G_k(t, s)| ds$ . Letting  $M_4 = \frac{3}{2}N_1N_2 + D_1N_2 + \frac{3}{2}M_2M_3 + D_2M_3 + M_1$ ,  $M_5 = \frac{N_1}{2}N_2 + \frac{M_2}{2}M_3$ ,  $\lambda = \lambda_1 + \lambda_2$  and solving algebraically for  $\|p_{n+1}\|$ , we obtain

$$\|p_{n+1}\| \leq \frac{1}{1 - \lambda} [M_4\|p_n\|^2 + M_5\|q_n\|^2].$$

#### 4. Concluding Remarks

We have developed the generalized quasilinearization method for a forced Duffing equation with mixed three-point nonlinear boundary conditions. Several interesting results can be recorded as a special case of the work established in this paper, for example, if we take  $g_1(x(\frac{1}{2})) = a$  and  $g_2(x(\frac{1}{2})) = b$  ( $a$  and  $b$  are constants), the problem corresponds to a two-point problem involving a forced

Duffing equation with separated boundary conditions. The classical method of quasilinearization for a forced Duffing equation with three-point nonlinear boundary conditions can be recorded by taking  $\phi(t, x) \equiv 0$  and  $\psi_i(x) \equiv 0$  in the assumptions  $(A_2)$  and  $(A_3)$  of Theorem 3.

### References

- [1] B. Ahmad, A quasilinearization method for a class of integro-differential equations with mixed nonlinearities, *Nonlinear Analysis, Real World Appl.*, In Press.
- [2] B. Ahmad, Monotone iteration scheme for general three-point nonlinear boundary value problems, *New Zealand J. Math.*, To Appear.
- [3] B. Ahmad, A. Al-Saedi, S. Sivasundaram, Approximation of the solution of nonlinear second order integro-differential equations, *Dynamic Systems Appl.*, **14** (2005), 253-263.
- [4] B. Ahmad, J.J. Nieto, N. Shahzad, The Bellman-Kalaba-Lakshamikantham quasilinearization method for Neumann problems, *J. Math. Anal. Appl.*, **257** (2001), 356-363.
- [5] C. Bai, J. Fang, Existence of multiple positive solutions for nonlinear m-point boundary value problems, *J. Math. Anal. Appl.*, **281** (2003), 76-85.
- [6] R. Bellman, *Methods of Nonlinear Analysis*, Volume 2, Academic Press, New York (1973).
- [7] R. Bellman, R. Kalaba, *Quasilinearization and Nonlinear Boundary Value Problems*, Amer. Elsevier, New York (1965).
- [8] A. Buica, Quasilinearization for the forced Duffing equation, *Studia Uni. Babeş-Bolyai Math.*, **47** (2000), 21-29.
- [9] A. Cabada, J.J. Nieto, Quasilinearization and rate of convergence for higher order nonlinear periodic boundary value problems, *J. Optim. Theory Appl.*, **108** (2001), 97-107.
- [10] W. Coppel, Disconjugacy, *Lecture Notes in Mathematics*, Volume **220**, Springer-Verlag, New York-Berlin (1971).

- [11] P.W. Eloe, B. Ahmad, Positive solutions of a nonlinear  $n$ -th order boundary value problem with nonlocal conditions, *Appl. Math. Letters*, **18**, No. 5 (2005), 521-527.
- [12] P. Eloe, Y. Gao, The method of quasilinearization and a three-point boundary value problem, *J. Korean Math. Soc.*, **39**, No. 2 (2002), 319-330.
- [13] C.P. Gupta, A second order  $m$ -point boundary value problem at resonance, *Nonlinear Anal.*, **24** (1995), 1483-1489.
- [14] C.P. Gupta, S. Trofimchuck, A priori estimates for the existence of a solution for a multi-point boundary value problem, *J. Inequal. Appl.*, **5** (2000), 351-365.
- [15] L. Jackson, Boundary value problems for ordinary differential equations, In: *Studies in Ordinary Differential Equations* (Ed. J.K. Hale), MAA Studies in Mathematics, Mathematical Association of America, **14** (1977), 93-127.
- [16] I.T. Kiguradze, A.G. Lomtadze, On certain boundary value problems for second-order linear ordinary differential equations with singularities, *J. Math. Anal. Appl.*, **101** (1984), 325-347.
- [17] V. Lakshmikantham, An extension of the method of quasilinearization, *J. Optim. Theory Appl.*, **82** (1994), 315-321.
- [18] V. Lakshmikantham, Further improvement of generalized quasilinearization, *Nonlinear Anal.*, **27** (1996), 223-227.
- [19] V. Lakshmikantham, A.S. Vatsala, *Generalized Quasilinearization for Nonlinear Problems*, Kluwer Academic Publishers, Dordrecht (1998).
- [20] E.S. Lee, *Quasilinearization and Invariant Embedding*, Academic Press, New York (1968).
- [21] R. Ma, Existence theorems for for a second order three-point boundary value problem, *J. Math. Anal. Appl.*, **212** (1997), 430-442.
- [22] R. Ma, Existence and uniqueness of solutions to first-order three-point boundary value problems, *Appl. Math. Lett.*, **15** (2002), 211-216.
- [23] V.B. Mandelzweig, F. Tabakin, Quasilinearization approach to nonlinear problems in physics with application to nonlinear ODEs, *Computer Physics Comm.*, **141** (2001), 268-281.

- [24] J.J. Nieto, A. Torres, A nonlinear biomathematical model for the study of intracranial aneurysms, *J. Neurological Science*, **177** (2000), 18-23.
- [25] S. Nikolov, S. Stoytchev, A. Torres, J.J. Nieto, Biomathematical modeling and analysis of blood flow in an intracranial aneurysms, *Neurological Research*, **25** (2003), 497-504.
- [26] I. Yermachenko, F. Sadyrbaev, Quasilinearization and multiple solutions of the Emden-Fowler type equation, *Math. Model. Anal.*, **10** (2005), 41-50.