

CHAIN CODES AND SPHERICAL TITS BUILDINGS

Luciano Panek^{1 §}, Marcelo Firer²

¹State University of the West of the Paraná

Av. Tarquínio Joslin dos Santos 1300

Foz do Iguaçu, 85870-650, PR, BRAZIL

e-mail: lucpanek@gmail.com

²State University of Campinas

Cx. Postal 6065, Campinas, 13081-970, SP, BRAZIL

e-mail: mfirer@ime.unicamp.br

Abstract: To an n -dimensional vector space V over a finite field \mathbf{F}_q there is an (naturally) associated spherical building of type A_{n-1} . The chambers of such a building are maximal flags: maximal sequences of nested subspaces of V . In the case $q = 2$, there is a unique $(n - 1)$ -dimensional maximum distance separable code in V . We show the existence of chambers associated to such a code that are of chain type (in the sense of coding theory) and give a complete characterization of the connected components of the chain type chambers.

AMS Subject Classification: 94B05, 14M15, 14NXX

Key Words: Hamming weights, chain codes, spherical Tits buildings

1. Introduction

In 1991 Wei introduced the concept of generalized minimum Hamming weights (see [6]) motivated by several applications in cryptography. With different motivation, similar properties of irreducible cyclic codes were studied by Hellesteth, Kløve and Mykkeltveit in 1977 (see [1]).

Since generalized weights were introduced several results were obtained generalizing already known results in coding theory. A major part of these results

Received: December 12, 2006

© 2007, Academic Publications Ltd.

§Correspondence author

can be found in the work [4] of Tsfasman and Vlăduț where the generalized weights are calculated through the projective systems (a set of points in a projective space over a finite field; see [5]).

Throughout this paper all codes are linear and defined over the finite field \mathbf{F}_2 of order two. If D is a subcode of the linear code C we write $D \leq C$ and $D < C$ in case it is a proper subspace of C . The r -th *minimum Hamming weight* of a k -dimensional linear code $C \leq \mathbf{F}_2^n$, is defined as

$$d_r(C) = \min \{ \|D\| : D \leq C, \dim(D) = r \},$$

with

$$\|D\| = \# \bigcup_{v \in D} \text{Supp}(v),$$

where $\text{Supp}(v)$, the *support* of $v = (v_1, \dots, v_n)$, is $\{i : v_i \neq 0\}$. A code C with *minimal weights hierarchy* $(\delta_1, \dots, \delta_k)$, where $\delta_r := d_r(C)$, is called an $[n; k; \delta_1, \dots, \delta_k]_2$ -code.

As a consequence of the definition we have that

$$1 \leq d_1(C) < d_2(C) < \dots < d_k(C) \leq n$$

(see [6, Theorem 1]) and it follows that

$$r \leq d_r(C) \leq n - k + r$$

for all $r \in \{1, \dots, k\}$ (see [6, Theorem 10]). A code such that $d_r(C) = n - k + r$ for some given $r \in \{1, \dots, k\}$ is called r -MDS code (r -maximum distance separable). If C is a r -MDS code, then it is also s -MDS for any $s \geq r$.

In this work we will study a special family of codes in \mathbf{F}_2^n , introduced by Wei and Yang [7], called chain codes. A code C is called a *chain code* (or *code of chain type*) if there is a sequence of sub-codes

$$D_1 < D_2 < \dots < D_k = C,$$

with $\|D_r\| = d_r(C)$ for every $r = 1, 2, \dots, k$, where $k = \dim(C)$. This is a particular but significant class of codes, since it includes the Hamming and the dual Hamming codes, Reed-Muller codes of any order, 1-MDS codes and the Golay codes (see [7, Theorem 6]). We observe that we can describe a maximal flag by giving an ordered base $\{v_1, \dots, v_k\}$ such that for every $i = 1, 2, \dots, k$, D_i is generated by $\{v_1, \dots, v_i\}$. Similar construction can be made for other flags, not necessarily maximal.

The main result of this work is Theorem 17, where we show that the set of flags (over the field \mathbf{F}_2) associated to a chain code of codimension 1 with

fixed weight hierarchy is connected, in the sense there is a sequence of flags $\alpha = \alpha_1, \alpha_2, \dots, \alpha_m = \beta$, where all α_i has the same weight hierarchy and each (α_i, α_{i+1}) can be determined by ordered bases that differ only by a single transposition.

This result is obtained through the characterization of the maximal flags associated to chain codes as chambers of a spherical Tits building, as we will see in the next section. This characterization, together with Theorem 17, enable us to count all the maximal flags with a given weight hierarchy (Proposition 16 and Corollary 14).

In Section 2 we introduce basic concepts related to Tits buildings and enunciate the proposed problems in terms of this structure. In Section 3 we study in details the structure of chain codes with weights hierarchy $(2, 3, \dots, n)$, beginning the counting procedures, mainly Theorem 5. Finally, in Section 4 we present the main results in this work: considering the family of all flags associated to chain codes of codimension 1, we have that each connected component of this family is determined exclusively by the weights hierarchies $(1, 2, \dots, n - 1)$ and $(2, 3, \dots, n)$ (Theorem 17 and Corollary 18).

2. Spherical Tits Buildings

We begin this section with generic definitions on abstract chamber systems and basic concepts of Tits buildings. We present only the concepts that are strictly necessary for this work, refereing the reader to [3] for more details.

A *chamber system* over a set I is triple $(\Lambda, \overset{i}{-}, I)$, where Λ is a set, I is a set of indices and for each $i \in I$, $\overset{i}{-}$ is an equivalence relation in Λ . The elements of Λ are called *chambers* and if $\alpha \overset{i}{-} \beta$ we say that the chambers α and β are *i -adjacent*, or simply adjacent if we do not need to distinguish the adjacency type. A *gallery of length k and type $i_1 i_2 \dots i_k$* joining two chambers α and β is a finite sequence of chambers $\alpha = \alpha_0, \alpha_1, \dots, \alpha_k = \beta$ such that the chambers α_{j-1} and α_j are different but i_j -adjacent for every $j \in \{1, \dots, k\}$. A *minimal gallery* joining α and β is a (not necessarily unique) gallery of minimal length joining the chambers.

If every i_j belongs to some subset $J \subset I$, we say the gallery is a *J -gallery*. A chamber system Λ , or a subsystem $\Lambda' \subseteq \Lambda$ is called *connected* (*J -connected*) if any two chambers can be connected (joined) by a gallery (*J -gallery*). In this case, we define the *distance* $d(\alpha, \beta)$ between two chambers as the length of a *minimal gallery* joining the chambers. A subset $\Lambda' \subseteq \Lambda$ is called *convex* if every

minimal gallery between any two chambers of Λ' is entirely contained in Λ' .

The J -connected components are called *residues of type J* , or simply J -*residues*. We denote the equivalence class (J -residue) of an element $\alpha \in \Lambda$ by $\text{Res}(\alpha; J)$. The *rank* of chamber system over set I is the cardinality of I , and the *corank* of $J \subset I$ is the cardinality of $I \setminus J$. A *morphism* $\phi : \Lambda \rightarrow \Gamma$ between two chamber systems over the same index set I is a map defined on the chambers that preserves i -adjacency for every $i \in I$. An *isomorphism* is an invertible morphism such that the inverse is also a morphism.

A *Coxeter group* is a group W finitely generated by a set $\{r_1, \dots, r_n\}$, subject only to the relations $(r_i r_j)^{m_{ij}} = 1$, where $m_{ij} \in \mathbf{N} \cup \{\infty\}$, $m_{ii} = 1$, and $m_{ij} = m_{ji}$ for every $i, j \in I = \{1, \dots, n\}$. The matrix $(m_{ij})_{i,j=1}^n$ is called the *Coxeter matrix* of W and denoted for M_I . The generators of W define a structure of chamber systems over I in the group in a canonical way: two elements $w, w' \in W$ are said to be i -adjacent if and only if $w' = w r_i$. This system is called a *Coxeter complex*, the fundamental “bricks” that constitute a Tits building.

Definition 1. Let Δ be a chamber system and Σ a family of subsystems, all isomorphic to a given finite Coxeter complex, such that:

- (i) For any two chambers there is $\Sigma \in \Sigma$ containing both of them;
- (ii) For each pair $\Sigma, \Sigma' \in \Sigma$ with a chamber in common there is an isomorphism of chamber systems $\phi : \Sigma \rightarrow \Sigma'$ that fixes $\Sigma \cap \Sigma'$ pointwise;

Then the pair (Δ, Σ) is called a *spherical Tits buildings* and the subsystems of Σ *apartments*.

Since a Coxeter complex is a finite complex system, it can be realized as a complex structure on a metric sphere (see [2]).

Example 2. Let \mathbf{F}_q be a finite field with q elements and \mathbf{F}_q^n the vector space of dimension n . We use the notation $(D_i)_{i=1}^{l-1}$ to represent a *flag of length $l-1$*

$$\{0\} = D_0 \leq D_1 \leq D_2 \leq \dots \leq D_{l-1} \leq D_l = \mathbf{F}_q^n.$$

A *maximal flag* is a flag of length n , and in this case $\dim(D_j) = j$. If we consider the building (Δ, Σ) defined as follows. The chambers of the building would be the set of all maximal flags:

$$\Delta = A_{n-1}(q) = \left\{ (D_i)_{i=1}^{n-1} : \dim(D_j) = j, D_j \subset \mathbf{F}_q^n \right\}.$$

Given a set $\{u_1, \dots, u_m\}$ of linearly independent vectors, we denote by

$$((u_1, \dots, u_i)_{i=1}^m)$$

the maximal flag $(D_i)_{i=1}^m$ such that D_i is spanned by $\{u_1, \dots, u_i\}$. Given a base $\{v_1, \dots, v_n\}$ of the linear space \mathbf{F}_q^n , the set of all flags obtained by permutations of this base is an apartment

$$\Sigma = \left\{ \left(\langle v_{\sigma(1)}, \dots, v_{\sigma(i)} \rangle \right)_{i=1}^{n-1} : \sigma \in \mathbf{S}_n \right\}$$

and

$$\Sigma = \left\{ \left\{ \left(\langle v_{\sigma(1)}, \dots, v_{\sigma(n)} \rangle \right)_{i=1}^{n-1} : \sigma \in \mathbf{S}_n \right\} : \{v_1, \dots, v_n\} \text{ is a base of } \mathbf{F}_q^n \right\},$$

where \mathbf{S}_n is the symmetric group. We note that different bases can give rise to the same chamber, or even the same apartment. Two chambers $(D_i)_{i=1}^{n-1}$ and $(D'_i)_{i=1}^{n-1}$ in Δ are i -adjacent if $D_j = D'_j$ for any $j \neq i$. The Coxeter group associated to the building $A_{n-1}(q)$ is the symmetric group \mathbf{S}_n .

Example 3. Every J -residue of a Tits building Δ of type M_I is a Tits building of type M_J (see [3, Theorem 3.5]).

3. Connecting Chambers by Galleries

We denote by $\Delta_2(\delta_1, \delta_2, \dots, \delta_{n-1})$ the set of all maximal flags $D_1 < \dots < D_{n-1}$ of \mathbf{F}_2^n such that

$$\|D_i\| = \delta_i := d_i(D_{n-1})$$

for all $i \in \{1, 2, \dots, n-1\}$. In other words, $\Delta_2(\delta_1, \delta_2, \dots, \delta_{n-1})$ is the set of all maximal flags of \mathbf{F}_2^n that achieve the weights hierarchy $(\delta_1, \delta_2, \dots, \delta_{n-1})$. Each of those flags is called a *chambers of type* $(\delta_1, \delta_2, \dots, \delta_{n-1})$.

We denote by $M^{n-1}(2) < \mathbf{F}_2^n$ the unique maximal non trivial 1-MDS code of \mathbf{F}_2^n . This code is the $(n-1)$ -dimensional subspace of \mathbf{F}_2^n consisting of all vectors with even weight. So, if α is a chamber (flag) of type $(2, 3, \dots, n)$ such that $M^{n-1}(2)$ is its maximal non-trivial subspace, we have that

$$\Delta_2(2, 3, \dots, n) \subset \text{Res}(\alpha, \{1, 2, \dots, n-2\}).$$

Given $\alpha \in \Delta_2(2, 3, \dots, n)$, the j -sphere with center α and ray 1 is the set of chambers $\beta \neq \alpha$ in $\Delta_2(2, 3, \dots, n)$ j -adjacent to α :

$$B_j(\alpha) = \left\{ \beta \in \Delta_2(2, 3, \dots, n) : d(\alpha, \beta) = 1 \text{ and } \beta \stackrel{j}{\sim} \alpha \right\}.$$

We will show that those stars are rather trivial. We start with a lemma.

Lemma 4. *Let $C \leq \mathbf{F}_q^n$ and*

$$E_i(C) = \# \{D : D \leq C, \dim(D) = 1, \|D\| = i\} .$$

Given $j \in \{1, \dots, n - 1\}$ consider $D_j \leq M^{n-1}(2)$ such that $\dim(D_j) = j$ and $\|D_j\| = j + 1$. Then

$$E_i(D_j) = \begin{cases} 0 & \text{if } i \text{ is odd,} \\ \binom{j+1}{i} & \text{if } i \text{ is even.} \end{cases}$$

Proof. Let $\{m_1, \dots, m_{j+1}\} = \text{Supp}(D_j)$. If $\pi : \mathbf{F}_2^n \rightarrow \mathbf{F}_2^{j+1}$ is the projection $\pi(x_1, \dots, x_n) = (x_{m_1}, \dots, x_{m_{j+1}})$, then $\pi(D_j) \leq \mathbf{F}_2^{j+1}$ is a subcode of dimension j . Since $D_j \leq M^{n-1}(2)$, we find that the Hamming weight of any word in $\pi(D_j)$ is even. Then

$$\pi(D_j) = M^{j+1}(2),$$

and the result follows. □

Let $U \leq V \leq \mathbf{F}_2^n$ be linear subspaces, with dimensions r and t respectively. The number of s -dimensional subspaces of V containing U equals (see [4, Lemma 2.2])

$$\begin{bmatrix} t - r \\ s - r \end{bmatrix} := \prod_{i=1}^{s-r} \frac{(2^{t-r} - 2^{i-1})}{(2^{s-r} - 2^{i-1})}.$$

Theorem 5. *Let $n > 2$ and $\alpha \in \Delta_2(2, 3, \dots, n)$. Then*

$$\#B_i(\alpha) = \begin{cases} 2 & \text{if } i = 1, \\ 1 & \text{if } i \neq 1. \end{cases}$$

In particular, $B_1(\alpha) \cup \{\alpha\} = \text{Res}(\alpha; \{1\})$.

Proof. Let $\alpha := (D_i)_{i=1}^{n-1} \in \Delta_2(2, 3, \dots, n)$. Since $\dim(D_{i+1}/D_{i-1})=2$, it has exactly 3 subspaces of dimension 1. One of them corresponds to the projection of D_i and the other two define the two chambers i -adjacent to $(D_i)_{i=1}^{n-1}$, and those are not necessarily contained in $\Delta_2(2, 3, \dots, n)$.

If $i = 1$, it is enough to notice that for any $v \in D_2 \setminus \{0\}$, $\|v\| = 2$ and this means that every chamber 1-adjacent to $(D_i)_{i=1}^{n-1}$ is in $\Delta_2(2, 3, \dots, n)$ and it follows that $\#B_1(\alpha) = 2$.

If $i \neq 1$, we consider $\{v_1, \dots, v_{i+1}\}$ a base of D_{i+1} such that $\{v_1, \dots, v_{i-1}\}$ generates D_{i-1} and $\{v_1, \dots, v_i\}$ generates D_i . Since D_{i+1} is a chain code, we can assume that

$$\# \left(\bigcup_{j=1}^l \text{Supp}(v_j) \right) = l + 1,$$

for every $l \in \{1, \dots, i + 1\}$. From Lemma 4, we find that D_i and D_{i+1} have respectively $\binom{i+1}{2}$ and $\binom{i+2}{2}$ words with weight 2. We claim there is $w_i \in D_{i+1}, w_i \neq v_i$ such that $\|w_i\| = 2$ and

$$\# \left(\left(\bigcup_{j=1}^{i-1} \text{Supp}(v_j) \right) \cup \text{Supp}(w_i) \right) = i + 1.$$

In fact, if $\text{Supp}(\langle v_1, \dots, v_{i+1} \rangle) = \{m_1, \dots, m_{i+2}\}$, there are $m_s \in \text{Supp}(v_{i+1})$ and $m_r \in \text{Supp}(\langle v_1, \dots, v_i \rangle)$ such that $m_s \notin \text{Supp}(\langle v_1, \dots, v_i \rangle)$ and then, every $w_i \in D_{i+1}$ such that $\text{Supp}(w_i) = \{m_r, m_s\}$ satisfies the desired conditions. So, we obtained two i -dimensional codes of chain type, let us say D_i and D'_i . The amount of distinct words of weight 2 in those codes equals

$$E_2(D_i) + E_2(D'_i) - E_2(D_{i-1}).$$

Since the number of words of weight 2 in $D_{i+1} \setminus (D_i \cup D'_i)$ is smaller than the number of words of weight 2 in D_i (or D'_i) that are not in D_{i-1} , that is,

$$E_2(D_{i+1}) - (E_2(D_i) + E_2(D'_i) - E_2(D_{i-1})) < E_2(D_i) - E_2(D_{i-1}),$$

we conclude that just two i -adjacent chambers can be of chain type, because the number of distinct words of weight 2 in a code $D \leq M^{n-1}(2)$ such that $\dim D = i$ and $\|D\| = i + 1$ equals $\binom{i+1}{2}$ (Lemma 4). \square

From here on, we will leave the trivial cases and consider only codes in n -dimensional spaces with $n > 2$.

If $B(\alpha) = \bigcup_j B_j(\alpha)$, then $\#B(\alpha)$ is called the *valency* of α , that is, the number of chambers in $\Delta_2(2, 3, \dots, n)$ adjacent to α . The next corollaries follows trivially from the preceding theorem.

Corollary 6. *Let $\alpha = (D_i)_{i=1}^{n-1} \in \Delta_2(2, 3, \dots, n)$. Then*

$$\# \left(\bigcup_{j=1}^r B_j(\alpha) \right) = \begin{cases} r + 1 & \text{if } r = 1, 2, \dots, n - 2, \\ r & \text{if } r = n - 1. \end{cases}$$

Corollary 7. *Let $\alpha \in \Delta_2(2, 3, \dots, n)$. Then the valency of α is $n - 1$.*

Suppose $D_i < D_{i+1} < M^{n-1}(2)$ are codes with $\|D_{i+1}\| = i + 2$. Since $D_i < D_{i+1}$ we have that $\|D_i\| \leq \|D_{i+1}\| = i + 2$. Since D_i is i -dimensional we cannot have $\|D_i\| \leq i$, because this means $\|D_i\| < d_i(M^{n-1}(2))$, contradicting the minimality of $d_i(M^{n-1}(2))$. It follows that $\|D_i\| = i + 1$ or $\|D_i\| = i + 2$.

Corollary 8. *Let*

$$\alpha \in \Delta_2(2, 3, \dots, n)$$

and

$$i \in \{2, 3, \dots, n - 2\}.$$

Then there is a unique chamber

$$(D_1, \dots, D_{n-1}) \in A_{n-1}(2)$$

that is i -adjacent to α , and such that $\|D_i\| = i + 2$.

Proof. We already know that for $i \geq 2$ there are only 2 chambers i -adjacent to α , exactly one of them of type $(2, 3, \dots, n)$ (Theorem 5). Since


$$\|D_i\| \in \{i + 1, i + 2\},$$

we must have one of them satisfying the equation $\|D_i\| = i + 2$. □

Despite the fact it is very simple, Corollary 8 has an interesting consequence: given two chambers

$$(D_1^1, \dots, D_{n-1}^1), (D_1^2, \dots, D_{n-1}^2) \in A_{n-1}(q)$$

such that $\dim(D_i^1 \cap D_i^2) = i - 1$ for any $i \in \{2, 3, \dots, n - 1\}$, they can be connected by a galleries of length $2n - 3$, as indicated in the following diagram (the indexes under the lines indicate the adjacency type):



If the initial and final chambers of the gallery above are in $\Delta_2(2, 3, \dots, n)$, the whole gallery is contained in $\Delta_2(2, 3, \dots, n)$, as follows from the next theorem.

Theorem 9. *If $(D_i^1)_{i=1}^{n-1}, (D_i^2)_{i=1}^{n-1} \in \Delta_2(2, 3, \dots, n)$ and*

$$\dim(D_i^1 \cap D_i^2) = i - 1$$

for any $i \in \{2, 3, \dots, n - 2\}$, then

$$\|D_r^1 \cap D_r^2\| = r$$

for every $r \in \{2, 3, \dots, n - 2\}$.

Proof. The proof is by induction on r . For $r = 2$, the result is trivial since every $0 \neq u \in D_2^i$ satisfies $\|u\| = 2$, for $i = 1, 2$.

We assume now that $\|D_i^1 \cap D_i^2\| = i$ for any $i \in \{3, \dots, r\}$ and suppose that $\|D_{r+1}^1 \cap D_{r+1}^2\| \neq r + 1$. By Corollary 8 we get that $\|D_{r+1}^1 \cap D_{r+1}^2\| = r + 2$. Let $\{i_1, \dots, i_r\}$ be the support of $D_r^1 \cap D_r^2$. Since

$$(D_r^1 \cap D_r^2) < (D_{r+1}^1 \cap D_{r+1}^2) < D_{r+1}^j \quad (j = 1, 2),$$

we have that

$$\text{Supp}(D_{r+1}^1 \cap D_{r+1}^2) = \{i_1, \dots, i_r, l_1, l_2\}$$

and since $(D_r^1 \cap D_r^2) < D_{r+1}^j$ ($j = 1, 2$) we have that

$$\text{Supp}(D_{r+1}^j) = \{i_1, \dots, i_r, i_{r+1}^j, i_{r+2}^j\} \quad (j = 1, 2).$$

But $(D_{r+1}^1 \cap D_{r+1}^2) < D_{r+1}^j$ ($j = 1, 2$), and we can assume with no loss of generality that $l_1 = i_{r+1}^j$ and $l_2 = i_{r+2}^j$. But then, $\text{Supp}(D_{r+1}^1) = \text{Supp}(D_{r+1}^2)$, and since $\|D_{r+1}^1\| = \|D_{r+1}^2\| = j + 1$, it follows from Lemma 4 that $D_{r+1}^1 = D_{r+1}^2$, and $\dim(D_{r+1}^1 \cap D_{r+1}^2) = r + 1$, contradicting the hypothesis that $\dim(D_{r+1}^1 \cap D_{r+1}^2) = r$. \square

Corollary 10. *If $(D_i^1)_{i=1}^{n-1}, (D_i^2)_{i=1}^{n-1} \in \Delta_2(2, 3, \dots, n)$, $\dim(D_i^1 \cap D_i^2) = i - 1$ for every $i \in \{2, 3, \dots, n - 3\}$ and $\#(\text{Supp}(D_1^1) \cap \text{Supp}(D_1^2)) = 1$, then*

$$\|D_r^1 + D_r^2\| = r + 2$$

for every $r \in \{1, 2, \dots, n - 3\}$.

Proof. From Theorem 9 we know that $\|D_i^1 \cap D_i^2\| = i$ for any $i \in \{2, 3, \dots, n - 3\}$. Since $\|D_i^1 + D_i^2\| = \|D_i^1\| + \|D_i^2\| - \|D_i^1 \cap D_i^2\|$, we have that $\|D_i^1 + D_i^2\| = i + 2$ for any $i \in \{2, 3, \dots, n - 3\}$. For the case $i = 1$, we notice that $\#\{\text{Supp}(D_1^1) \cap \text{Supp}(D_1^2)\} = 1$, and $\|D_1^1\| = \|D_1^2\| = 2$ and find that $\|D_1^1 + D_1^2\| = 3$. \square

4. Connected Components

The main result of this work is presented in Theorem 17, where we characterize the connected components of maximal flags associated to the chain codes. To prove that $\Delta_2(2, 3, \dots, n)$ is connected, we will show that the set of chambers of type $(2, 3, \dots, n)$ can be described (see Example 3 for details) as the disjoint union of apartments in the Tits building

$$\text{Res}(\alpha; \{1, 2, \dots, n - 2\}), \alpha \in \Delta_2(2, 3, \dots, n).$$

To understand the structure of those buildings defined by residues, we take a close look at the 4-dimensional case. We notice that $\Delta_2(2, 3, 4)$ is the union of the four apartments defined by the bases bellow:

$$\begin{aligned} & \{(1, 0, 0, \underline{1}), (0, 1, 0, \underline{1}), (0, 0, 1, \underline{1})\}, \{(0, \underline{1}, 0, 1), (0, \underline{1}, 1, 0), (1, \underline{1}, 0, 0)\}, \\ & \{(0, 1, \underline{1}, 0), (1, 0, \underline{1}, 0), (0, 0, \underline{1}, 1)\}, \{(\underline{1}, 0, 0, 1), (\underline{1}, 1, 0, 0), (\underline{1}, 0, 1, 0)\}. \end{aligned}$$

The underlined coordinates suggest the picture we wish to generalize: we chose a coordinate to be nonzero, which defines a co-dimension 1 subspace, and the vectors with weight two constitute a base of it. This will produce the apartments of $\Delta_2(2, 3, \dots, n)$ and constitute the foundation of the proof that $\Delta_2(2, 3, \dots, n)$ is connected.

Lemma 11. Given $(D_i)_{i=1}^{n-1} \in \Delta_2(2, 3, \dots, n)$ there is a base $\{v_1, \dots, v_{n-1}\}$ of $M^{n-1}(2)$ and $l \in \{1, 2, \dots, n\}$ such that:

- (i) $\langle v_1, \dots, v_r \rangle = D_r$, for any $r \in \{1, \dots, n-1\}$;
- (ii) $\text{Supp}(v_i) \cap \text{Supp}(v_j) = \{l\}$, for any $i, j \in \{1, \dots, n-1\}$ with $i \neq j$.

Proof. Since $M^{n-1}(2)$ is of chain type, there is a base $\{w_1, \dots, w_{n-1}\}$ of $M^{n-1}(2)$ such that

$$\# \left(\bigcup_{j=1}^i \text{Supp}(w_j) \right) = d_i(M^{n-1}(2))$$

and

$$\langle w_1, \dots, w_i \rangle = D_i,$$

with $i \in \{1, \dots, n-1\}$. We note that since $d_1(M^{n-1}(2)) = 2$ we have that $\|w_1\| = 2$, let us say $\text{Supp}(w_1) = \{i_1, i_2\}$ and $\text{Supp}(w_j) = \{i_j, i_{j+1}\}$, $j \in \{2, 3, \dots, n-1\}$, with $i_j \in \text{Supp}(w_{j-1})$ and

$$i_{j+1} \in \{1, \dots, n-1\} \setminus \text{Supp}(D_{j-1}).$$

Defining

$$\begin{aligned} v_1 &= w_1, \\ v_j &= v_{j-1} + w_j, \quad j = 2, \dots, n-1 \end{aligned}$$

we find that

$$\begin{aligned} \text{Supp}(v_1) &= \{i_1, i_2\}, \\ \text{Supp}(v_j) &= \{i_1, i_{j-1}\}, \quad j = 2, \dots, n-1. \end{aligned}$$

Then $\text{Supp}(v_i) \cap \text{Supp}(v_j) = \{i_1\}$ for any $i, j \in \{1, \dots, n-1\}$ with $i \neq j$. Therefore $\{v_1, \dots, v_{n-1}\}$ is a base satisfying the requested properties. \square

Let us define the vector $v_j^i \in \mathbf{F}_2^n$ as

$$v_j^i := \begin{cases} (0, \dots, 0, 1_i, 0, \dots, 0, 1_{j+1}, 0, \dots, 0) & \text{if } j \geq i, \\ (0, \dots, 0, 1_j, 0, \dots, 0, 1_i, 0, \dots, 0) & \text{if } j < i, \end{cases}$$

where the subindex indicate the corresponding coordinate we are assigning non-zero values. As $M^{n-1}(2)$ is an $(n-1)$ -dimensional subspace consisting of all the vectors of even weight of \mathbf{F}_2^n and $\{v_1^i, v_2^i, \dots, v_{n-1}^i\}$ is linearly independent,

we find that for each $i \in \{1, \dots, n\}$, the set $\{v_j^i : j = 1, \dots, n-1\}$ is a base of $M^{n-1}(2)$:

$$M^{n-1}(2) = \langle v_j^i : j = 1, \dots, n-1 \rangle.$$

So it defines the apartment $\Sigma_i \subset \Delta_2(2, 3, \dots, n)$:

$$\Sigma_i = \left\{ \left(\langle v_{\sigma(1)}^i, \dots, v_{\sigma(j)}^i \rangle \right)_{j=1}^{n-1} : \sigma \in \mathbf{S}_{n-1} \right\}.$$

Proposition 12. *With the notation above defined,*

$$\Delta_2(2, 3, \dots, n) = \bigcup_{i=1}^n \Sigma_i.$$

Proof. By construction, $\bigcup_{i=1}^n \Sigma_i \subset \Delta_2(2, 3, \dots, n)$ and it is left to prove that each chamber $\alpha = (D_i)_{i=1}^{n-1} \in \Delta_2(2, 3, \dots, n)$ is contained in some of those apartments. Let us consider a base $\{v_1, \dots, v_{n-1}\}$ of $M^{n-1}(2)$ such that $\langle v_1, \dots, v_r \rangle = D_r$ and $\text{Supp}(v_i) \cap \text{Supp}(v_j) = \{l\}$, whenever $i \neq j$ and $r \in \{1, 2, \dots, n-1\}$ (existence guaranteed by Lemma 11). Setting

$$i_1 \in \text{Supp}(v_1) \setminus \{l\}, \dots, i_{n-1} \in \text{Supp}(v_{n-1}) \setminus \{l\}$$

and

$$v^{i,j} = (0, \dots, 0, 1_i, 0, \dots, 0, 1_j, 0, \dots, 0),$$

we have that $\alpha = (D_i)_{i=1}^{n-1} = (\langle v^{l,i_1}, \dots, v^{l,i_j} \rangle)_{j=1}^{n-1} \in \Sigma_l$ and it follows that $\Delta_2(2, 3, \dots, n) = \bigcup_{i=1}^n \Sigma_i$. \square

Proposition 13. *Let $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$. Then $\#(\Sigma_i \cap \Sigma_j) = (n-2)!$.*

Proof. Consider two apartments Σ_i and Σ_j . Let $\sigma \in \mathbf{S}_{n-1}$ such that $\sigma(1) = j$, and suppose that $j < i$. Then

$$\left(\langle v_{\sigma(1)}^i, \dots, v_{\sigma(l)}^i \rangle \right)_{l=1}^{n-1} \in \Sigma_i.$$

Let $h \in \{2, 3, \dots, n\}$. Since

$$v_{\sigma(1)}^i + v_{\sigma(h)}^i = \begin{cases} v^{j, \sigma(h)} & \text{if } \sigma(h) \in \{1, 2, \dots, i-1\}, \\ v^{j, \sigma(h)+1} & \text{if } \sigma(h) \in \{i, i+1, \dots, n-1\}, \end{cases}$$

we find that

$$\left(\langle v_{\sigma(1)}^i, \dots, v_{\sigma(l)}^i \rangle \right)_{l=1}^{n-1} \in \Sigma_j.$$

and therefore $\Sigma_i \cap \Sigma_j \neq \emptyset$. Since we have a total of $(n-2)!$ permutations of type $(1j \dots) \in \mathbf{S}_{n-1}$, we conclude that $\#(\Sigma_i \cap \Sigma_j) = (n-2)!$. \square

Corollary 14. *The number of chambers in $\Delta_2(2, 3, \dots, n)$ is $n!/2$.*

Proof. We note that

$$(\Sigma_l \cap \Sigma_i) \cap (\Sigma_l \cap \Sigma_j) = \emptyset,$$

if $i \neq j$ and $i, j \neq l$. So, if we assume that $i \neq j$ with $i, j \neq l$, we have that

$$(\Sigma_l \cap \Sigma_1) \cup (\Sigma_l \cap \Sigma_2) \cup \dots \cup (\Sigma_l \cap \Sigma_{l-1})$$

is a disjoint union. The result follows from Principle of Inclusion-Exclusion:

$$\begin{aligned} \#\Delta_2(2, 3, \dots, n) &= \#(\Sigma_1 \cup \dots \cup \Sigma_n) \\ &= \#\Sigma_1 + \sum_{i=2}^n \left(\#\Sigma_i - \sum_{j=1}^{i-1} \#(\Sigma_i \cap \Sigma_j) \right) = (n-1)! + (n-2)(n-1)! \\ &\quad - \frac{(n-1)(n-2)}{2}(n-2)! = \frac{n!}{2}. \quad \square \end{aligned}$$

Corollary 15. *For any $n \geq 3$, $\Delta_2(2, 3, \dots, n)$ is connected.*

Proof. Let $\alpha, \beta \in \Delta_2(2, 3, \dots, n)$. If $\alpha, \beta \in \Sigma_i$ for some i , then the chambers can be connected by a gallery, since each apartment Σ_i is convex (see [3, Theorem 3.8]). Suppose now that $\alpha \in \Sigma_i$ and $\beta \in \Sigma_j$ with $\alpha, \beta \notin \Sigma_i \cap \Sigma_j$. Since the intersection is not empty (Proposition 13), we can connect α to $\gamma \in \Sigma_i \cap \Sigma_j$ through a gallery in Σ_i , and β to γ through a gallery in Σ_j . Making the juxtaposition of these two galleries, we obtain a gallery joining α to β (passing through γ), entirely contained in $\Sigma_i \cup \Sigma_j \subset \Delta_2(2, 3, \dots, n)$. \square

Proposition 16. *The set of the chambers $\Delta_2(1, 2, \dots, n-1)$ is an apartment of $A_{n-1}(2)$. In particular, $\Delta_2(1, 2, \dots, n-1)$ is convex.*

Proof. Consider the canonical base $\{e_1, e_2, \dots, e_n\}$ of \mathbf{F}_2^n . It follows immediately from the definition that

$$\left\{ \left(\langle e_{\sigma(1)}, \dots, e_{\sigma(i)} \rangle \right)_{i=1}^{n-1} : \sigma \in \mathbf{S}_n \right\} \subseteq \Delta_2(1, 2, \dots, n-1).$$

Given $\alpha = (D_1, \dots, D_{n-1}) \in \Delta_2(1, 2, \dots, n-1)$ we choose a base $\{v_1, \dots, v_{n-1}\}$ of D_{n-1} such that

$$\# \left(\bigcup_{j=1}^i \text{Supp}(v_j) \right) = d_i(D_{n-1})$$

for any $i \in \{1, \dots, n-1\}$. We have to prove there is a permutation $\sigma \in \mathbf{S}_n$ such that

$$\langle v_1, \dots, v_i \rangle = \langle e_{\sigma(1)}, \dots, e_{\sigma(i)} \rangle, \quad i = 1, 2, \dots, n,$$

that is, up to re-scaling by scalars, this ordered base is just a permutation of the canonical one. Since $d_1(D_{n-1}) = 1$, we find that $v_1 = e_j$ for some $j \in \{1, \dots, n-1\}$. Since $\|\langle v_1, v_2 \rangle\| = 2$, we have that $\text{Supp}(v_2) = \{j, l\}$ or $\text{Supp}(v_2) = \{l\}$ (assuming, without loss of generality, that $l > j$). In the second case we have that $v_2 = e_l$. In the first case, we find that

$$v_2 = (0, \dots, 0, 1, 0, \dots, 0, 1, 0, \dots, 0)$$

and it follows that $-e_j + v_2 = e_l$. Proceeding with this process in the same manner, we find that $\alpha \in \left\{ \left(\langle e_{\sigma(1)}, \dots, e_{\sigma(i)} \rangle \right)_{i=1}^{n-1} : \sigma \in \mathbf{S}_n \right\}$.

The convexity follow of the fact that apartments are convex (see [3, Theorem 3.8]). □

Let us notice now that there are many codes $C \subset \mathbf{F}_2^n$ of codimension 1 with weights hierarchy $(1, 2, \dots, \widehat{m+1}, \dots, n)$. Indeed, any such code may be described as the kernel of a linear functional $\varphi : \mathbf{F}_2^n \rightarrow \mathbf{F}_2$; the code $M^{n-1}(2)$ is the kernel of $\varphi(v_1, \dots, v_n) = v_1 + \dots + v_n$. In the general case, the code of type $(1, 2, \dots, \widehat{m+1}, \dots, n)$ is the kernel of

$$\varphi(v_1, \dots, v_n) = v_1 + \dots + \widehat{v_{i_1}} + \dots + \widehat{v_{i_m}} + \dots + v_n. \tag{1}$$

It follows there are

$$n + \binom{n}{2} + \binom{n}{3} + \dots + \binom{n}{n-2} = 2(2^{n-1} - 1) - n$$

codes of codimension 1 in \mathbf{F}_2^n that are neither of the type $(2, 3, \dots, n)$ nor $(1, 2, \dots, n-1)$.

We denote by $\Delta_2\{i_1, \dots, i_m\}$ the set of chambers with weights hierarchy $(1, 2, \dots, \widehat{m+1}, \dots, n)$ that contains a codimension 1 code defined as the kernel of the functional (1).

Theorem 17. *The set $\Delta_2(1, 2, \dots, \widehat{m+1}, \dots, n)$ is a disjoint union of the J -connected components $\Delta_2 I$, with $I = \{i_1, \dots, i_m\}$ and $J = \{1, \dots, \widehat{m}, \dots, n-2\}$. Consequently the set*

$$\bigcup_{(d_1, \dots, d_{n-1})} \Delta_2(d_1, \dots, d_{n-1})$$

has exactly $2^n - n$ connected components.

Proof. Each partial flag $(D_i)_{i=1}^m \in \Delta_2 \{i_1, \dots, i_m\}$ may be described as the sequence of subspaces defined by an ordered base $\{e_{i_1}, \dots, e_{i_m}\}$ of D_m . In others word, the set of all such partial flags is J' -connected, for $J' = \{1, 2, \dots, m - 1\}$.

The partial flags $(D_i)_{i=m+1}^{n-1}$ in $\Delta_2 \{i_1, \dots, i_m\}$ are obtained from the vectors

$$(0, \dots, 0, 1_{j_1}, 0, \dots, 0, 1_{j_2}, 0, \dots, 0),$$

where j_1 is the first non-zero position in $\{1, 2, \dots, n\} \setminus I$ and $j_2 \in \{1, 2, \dots, n\} \setminus I$, $j_2 \neq j_1$. The set of all such partial flags is J'' -connected, $J'' = \{m + 1, m + 2, \dots, n - 2\}$. It follows that $\Delta_2 \{i_1, \dots, i_m\}$ is $J' \cup J''$ -connected.

Finally, we prove that

$$\Delta_2 \{i_1, \dots, i_m\} \cup \Delta_2 \{j_1, \dots, j_m\},$$

is not connected, except if one of those sets of indices is contained in the other. If we suppose neither one is contained in the other, there is $j_r \notin \{i_1, \dots, i_m\}$. Let $\alpha = (D_i)_{i=1}^m \in \Delta_2 \{i_1, \dots, i_m\}$ and $\beta = (D_i)_{i=1}^m \in \Delta_2 \{j_1, \dots, j_m\}$, and suppose that α and β can be connected by a gallery in $\Delta_2 \{i_1, \dots, i_m\} \cup \Delta_2 \{j_1, \dots, j_m\}$. But adjacency in $A_{n-1}(q)$ is defined by permutations of the elements of a given base, so that a gallery joining α to β needs to change, at some place m , the subspace D_m by the subspace D'_m . But in order to do so, we must have $\{e_{i_1}, \dots, e_{i_m}, e_{j_r}\} \subset D_{n-1} \cap D'_{n-1}$. But in this case, the subspace $\langle e_{i_1}, \dots, e_{i_m}, e_{j_r} \rangle \subset \mathbf{F}_2^n$ has dimension $m + 1$ and generalized weight equal $m + 1$, contradicting the minimality of $d_{m+1} = m + 2$. \square

Let $1 \leq r_1 < \dots < r_k \leq n$ be a sequence of integers, $I = \{i_1, \dots, i_m\}$, $N = \{1, 2, \dots, n\}$, $I \subset N$ and $I^c = N \setminus I$. We denoted by $\mathbf{F}^n(r_1, \dots, r_k)$ the set of all the flags $D_{r_1} < \dots < D_{r_k}$ formed by subspaces of \mathbf{F}_2^n such that $\dim(D_{r_j}) = r_j$. We defined the inclusions

$$\begin{aligned} i_I &: \mathbf{F}_2^m \rightarrow \mathbf{F}_2^n, \\ \widehat{i}_I &: A_{m-1}(2) \rightarrow \mathbf{F}^n(1, \dots, m) \end{aligned}$$

and

$$\widehat{i}_I : A_{n-m-1}(2) \rightarrow \mathbf{F}^n(m + 1, \dots, n - 1),$$

respectively as

$$i_I(x_1, \dots, x_m) = (0, \dots, 0, (x_1)_{i_1}, 0, \dots, 0, (x_m)_{i_m}, 0, \dots, 0),$$

$$\widehat{i}_I(D_1 < \dots < D_m) = i_I(D_1) < \dots < i_I(D_m)$$

and

$$\begin{aligned} \widehat{i}_I(D_1 < \dots < D_{n-m-1}) &= i_{I^c}(D_1) \oplus i_I(\mathbf{F}_2^m) \\ &< \dots < i_{I^c}(D_{n-m-1}) \oplus i_I(\mathbf{F}_2^m). \end{aligned}$$

Given chamber systems Λ_1, Λ_2 over I_1, I_2 , the *direct product* $\Lambda_1 \times \Lambda_2$ is a chamber system over the disjoint union $I_1 \cup I_2$. Its chambers are the pairs (α_1, α_2) , with $\alpha_i \in \Lambda_i$, and (α_1, α_2) is said to be *i*-adjacent to (β_1, β_2) for $i \in I_t$ ($t = 1$ or 2) if $\alpha_j = \beta_j$ for $j \neq t$ and $\alpha_t \overset{i}{\sim} \beta_t$ in Λ_t .

Notice now that the direct product

$$\widehat{i}_I(\Delta_2(1, 2, \dots, m-1)) \times \widehat{i}_I(\Delta_2(2, 3, \dots, n-m))$$

is isomorphic to the set $\Delta_2 I$. So, if we place $\Delta = \Delta_2(1, 2, \dots, m-1)$ and $\Delta' = \Delta_2(2, 3, \dots, n-m)$, we have the *coproduct*

$$\prod_{k=1}^{\binom{n}{m}} \Delta \times \Delta' = \bigcup_I \widehat{i}_I(\Delta) \times \widehat{i}_I(\Delta')$$

and as a particular case of Theorem 17 we have the following corollary.

Corollary 18. *The set $\Delta_2(1, 2, \dots, \widehat{m+1}, \dots, n)$ is isomorphic to the coproduct*

$$\prod_{k=1}^{\binom{n}{m}} \Delta \times \Delta',$$

where the codes of dimension m and $n - m - 1$ in each product $\Delta \times \Delta'$ are identified with the codes $\langle \{e_{i_j}\}_{j=1}^m \rangle$ and $\langle \{v_{j_2}^{j_1}\} \rangle$, $j_1, j_2 \in I^c$. Consequently $\Delta_2(1, 2, \dots, \widehat{m+1}, \dots, n)$ has exactly $n!/2$ chambers.

We have characterized the connected components of the union of chambers of chain type in \mathbf{F}_2^n , $\bigcup_{(d_1, \dots, d_{n-1})} \Delta_2(d_1, \dots, d_{n-1})$, and determined the cardinality of each such connected component. Those results are summarized in the table below.

$\Delta_2(1, 2, \dots, n-1)$	$n!$	Proposition 16
$\Delta_2(2, 3, \dots, n)$	$n!/2$	Corollary 15
$\Delta_2\{i_1, \dots, i_m\}$	$m!(n-m)!/2$	Theorem 17
$\Delta_2(1, 2, \dots, \widehat{m+1}, \dots, n)$	$n!/2$	Corollary 18

References

- [1] T. Helleseeth, T. Kløve, J. Mykkeltveit, The weight distribution of irreducible cyclic codes with block lengths $n_1((q^l - 1)/n)$, *Discrete Mathematics*, **18** (1977), 179-211.
- [2] J. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, Volume **29**, Cambridge University Press (1990).
- [3] M. Ronan, *Lectures on Buildings*, Perspectives in Mathematics, Volume **7**, Academic Press (1989).
- [4] M.A. Tsfasman, S.G. Vlăduț, Geometric approach to higher weights, *IEEE Trans. Inform. Theory*, **41**, No. 6 (1995), 1564-1588.
- [5] M.A. Tsfasman, S.G. Vlăduț, *Algebraic-Geometric Codes*, Mathematics and its Applications, Soviet Series, **58**, Kluwer Academic Publishers (1991).
- [6] V.K. Wei, Generalized Hamming weights for linear code, *IEEE Trans. Inform. Theory*, **37**, No. 5 (1991), 1412-1418.
- [7] V.K. Wei, K. Yang, On the generalized Hamming weights for product code, *IEEE Trans. Inform. Theory*, **39**, No. 5 (1993), 1709-1713.

