

**A PARTICULAR SOLUTION OF A NONLINEAR
BLACK-SCHOLES PARTIAL DIFFERENTIAL EQUATION**

Joseph Eyang'an Esekon

Department of Statistics and Actuarial Science

Maseno University

P.O. Box 333, Maseno, KENYA

Abstract: We study a nonlinear Black-Scholes partial differential equation whose nonlinearity is as a result of transaction costs that lead to market illiquidity. After reducing the equation into a nonlinear parabolic porous medium type equation, we find that the assumption of a traveling wave profile to the porous medium type equation reduces it further to ordinary differential equations. Solutions to all these transformed equations together with the use of localizing boundary conditions facilitate a twice continuously differentiable solution to the nonlinear Black-scholes equation. We also find that the option is always more volatile compared to the stock.

AMS Subject Classification: 35K10, 35K55

Key Words: porous medium equation, particular solution, illiquid markets, transaction cost

1. Introduction

Two primary assumptions are used in formulating classical arbitrage pricing theory: *frictionless* and *competitive* markets. There are no transaction costs and restrictions on trade in a frictionless market. In a competitive market, a trader can buy or sell any quantity of a security without changing its price.

The notion of liquidity risk is introduced on relaxing the assumptions above.

The purpose of this paper is to obtain a particular solution of the nonlinear Black-Scholes equation arising from transaction costs by Cetin *et al* in [2]. This is done by differentiating the equation twice with respect to the stock price. After substitutions and transformations, we get a nonlinear parabolic porous medium type equation. Assuming a traveling wave solution to the porous medium type equation reduces it further to ordinary differential equations (ODEs). All these transformed equations are solved to obtain a particular solution to the nonlinear Black-Scholes equation. A thorough analysis of this Black-Scholes equation is found in [3].

This paper is organized as follows. Section 2 describes the modified option pricing theory. The solution to the nonlinear Black-Scholes equation is presented in Section 3. Section 4 concludes the paper.

2. Modified Option Valuation Model

In this work, we will consider the continuous-time transaction-cost model for illiquid markets by Cetin *et al* in [2]. Two assets are used in the model: a bond (or a risk-free money market account with spot rate of interest $r \geq 0$) whose value at time t is $B_t \equiv 1$, and a stock. The stock is assumed to be risky and illiquid while the bond is assumed to be riskless and liquid.

In the transaction-cost model, a fundamental stock price process S_t^0 follows the dynamics

$$dS_t^0 = \mu S_t^0 dt + \sigma S_t^0 dW_t, \quad 0 \leq t \leq T,$$

where μ is drift, σ is volatility, T is time to expiry, and W_t is the Wiener process. When trading α shares, the transaction price to be paid at time t is

$$\bar{S}_t(\alpha) = e^{\rho\alpha} S_t^0, \quad \alpha \in \mathbb{R}, \quad (2.1)$$

where ρ is a liquidity parameter with $\rho \geq 0$. It is obvious from equation (2.1) that if $\rho > 0$ the trader will pay more than the fundamental price when buying the stock (i.e. $\alpha > 0$) and get less when selling (i.e. $\alpha < 0$). Therefore, a bid-ask-spread with size depending on α is essentially modeled by (2.1).

Taking the function $\phi(t, S_t^0)$ to be the number of shares traded rather than held and identifying this function to be a trading strategy, then for a Markovian trading strategy $\Phi_t = \phi(t, S_t^0)$ for a smooth function $\phi = u_S$, we have $\phi_S = u_{SS}$.

If the stock and bond positions are Φ_t and β_t respectively where Φ_t is a semimartingale, then the paper value is given by

$$V_t^M = \Phi_t S_t^0 + \beta_t.$$

The change in the quadratic variation,

$$\langle \Phi \rangle_t = \int_0^t (\phi_S(\tau, S_\tau^0) \sigma S_\tau^0)^2 d\tau,$$

is given by $d\langle \Phi \rangle_t = (u_{SS}(t, S_t^0) \sigma S_t^0)^2 dt$ since $\phi_S = u_{SS}$.

Applying Itô's formula to $u(t, S_t^0)$ gives

$$du(t, S_t^0) = u_S(t, S_t^0) dS_t^0 + (u_t(t, S_t^0) + \frac{1}{2} \sigma^2 (S_t^0)^2 u_{SS}(t, S_t^0)) dt. \tag{2.2}$$

Theorem A3 of [2] yields the wealth dynamics of a self-financing strategy as

$$\begin{aligned} dV_t^M &= \Phi_t dS_t^0 - \rho S_t^0 d\langle \Phi \rangle_t \\ &= u_S(t, S_t^0) dS_t^0 - \rho \sigma^2 (u_{SS}(t, S_t^0))^2 (S_t^0)^3 dt \end{aligned} \tag{2.3}$$

since $\Phi_t = \phi(t, S_t^0) = u_S$ and $d\langle \Phi \rangle_t = (u_{SS}(t, S_t^0) \sigma S_t^0)^2 dt$. Uniqueness of semimartingale decompositions means that the dt -terms in (2.2) and (2.3) must coincide. Equate these dt -terms. Applying a little algebra and then simplifying we get

$$u_t + \frac{1}{2} \sigma^2 S^2 u_{SS} (1 + 2\rho S u_{SS}) = 0, \quad u(S, T) = h(S_T), \tag{2.4}$$

where $h(S_T)$ is a terminal claim whose hedge cost, $u(S_t, t)$, is the solution to (2.4). For instance the terminal condition for a European call option is given by

$$u(S, T) = \max \{S - K, 0\} \quad \text{for } S \geq 0,$$

where $K > 0$ is the striking price. The boundary conditions for the option are as follows:

$$\begin{aligned} u(0, t) &= 0 \quad \text{for } 0 \leq t \leq T, \\ u(S, t) &\sim S - Ke^{-r(T-t)} \quad \text{as } S \rightarrow \infty. \end{aligned}$$

We take the last condition to mean that

$$\lim_{S \rightarrow \infty} \frac{u(S, t)}{S - Ke^{-r(T-t)}} = 1 \tag{2.5}$$

uniformly for $0 \leq t \leq T$.

The magnitude of the market impact is determined by ρS . Large ρ implies a big market impact of hedging. If $\rho = 0$, the asset's price follows the standard Black-Scholes model in [1] with constant volatility σ .

3. Solution to the Nonlinear Black-Scholes Equation

Lemma 3.1. *If $\nu(\xi)$ is a twice continuously differentiable positive function, and x and t are the spatial and time variables respectively, then there exists a traveling wave solution to the porous medium type equation,*

$$V_t + \frac{\sigma^2}{2}(VV_x + \frac{1}{2}V^2)_x = 0 \quad \text{in } \mathbb{R} \times [0, \infty), \quad (3.1)$$

of the form

$$V(x, t) = \nu(\xi) \quad \text{where } \xi = x - ct, \quad \xi \in \mathbb{R} \quad (3.2)$$

for $\sigma > 0$ such that $V = V(x, t)$ is a traveling wave of permanent form which translates to the right with constant speed $c > 0$.

Proof. Applying the chain rule to (3.2) gives

$$V_t = -c\nu'(\xi), \quad V_x = \nu'(\xi), \quad \text{and} \quad V_{xx} = \nu''(\xi),$$

where the prime denotes $\frac{d}{d\xi}$. Substituting these expressions into (3.1), we conclude that $\nu(\xi)$ must satisfy the nonlinear second order ODE

$$-c\nu' + \frac{\sigma^2}{2}(\nu\nu'' + (\nu')^2 + \nu\nu') = 0 \quad \text{in } \mathbb{R} \quad (3.3)$$

and hence V solves (3.1).

Assume also that the travelling wave solution $V(x, t)$ is *localized*. Then

$$\lim_{x \rightarrow \pm\infty} V(x, t) = \lim_{x \rightarrow \pm\infty} V_x(x, t) = \lim_{x \rightarrow \pm\infty} V_{xx}(x, t) = 0$$

or

$$\lim_{\xi \rightarrow \pm\infty} \nu(\xi) = \lim_{\xi \rightarrow \pm\infty} \nu'(\xi) = \lim_{\xi \rightarrow \pm\infty} \nu''(\xi) = 0. \quad (3.4)$$

We can now solve (3.3) in a closed-form by first writing it as

$$\frac{d}{d\xi} \left(\frac{\sigma^2}{2} \nu \nu' + \frac{\sigma^2}{4} \nu^2 - c\nu \right) = 0. \quad (3.5)$$

Integrating (3.5) and rearranging gives the standard form (see [4])

$$\nu' = \frac{2}{\sigma^2 \nu} (c\nu - \frac{\sigma^2}{4} \nu^2 + \kappa), \quad (3.6)$$

where κ is a constant of integration. From the localizing boundary conditions (3.4), $\kappa = 0$. Simplifying (3.6) further we realize that $\nu(\xi)$ satisfies the first order linear autonomous and separable ODE

$$-2 \frac{d\nu}{d\xi} = \nu - \frac{4c}{\sigma^2}.$$

Rearranging the equation above and integrating gives

$$\nu(\xi) = e^{\frac{\xi_0 - \xi}{2}} + \frac{4c}{\sigma^2}, \quad \sigma > 0, \tag{3.7}$$

where ξ_0 is a constant of integration. Hence,

$$V(x, t) = e^{\frac{x_0 - (x - ct)}{2}} + \frac{4c}{\sigma^2}, \quad \sigma > 0, \quad t \geq 0, \tag{3.8}$$

where x_0 is a constant of integration given by

$$\xi_0 = x_0 - c \cdot 0 = x_0$$

as $t = 0$. It is obvious from equations (3.7) and (3.8) that $\nu(\xi)$ and $V(x, t)$ are positive functions. □

Theorem 3.2. *If $V(x, t)$ is any positive solution to the porous medium type equation*

$$V_t + \frac{\sigma^2}{2} (VV_x + \frac{1}{2}V^2)_x = 0 \quad \text{in } \mathbb{R} \times [0, \infty),$$

then

$$u(S, t) = S - \frac{\sqrt{S_0}}{\rho} \left(\sqrt{S} e^{\frac{\sigma^2 t}{8}} + \frac{\sqrt{S_0}}{4} e^{\frac{\sigma^2 t}{4}} \right) \tag{3.9}$$

solves the nonlinear Black-Scholes equation

$$u_t + \frac{1}{2}\sigma^2 S^2 u_{SS} (1 + 2\rho S u_{SS}) = 0$$

for $S_0, S, \sigma, \rho > 0$ and $t \geq 0$, where S_0 is the initial stock price.

Proof. To obtain the solution to equation (2.4) we differentiate it twice with respect to S to get

$$w_t + \frac{\sigma^2 S^2}{2} (1 + 4\rho S w) w_{SS} + 2\rho\sigma^2 S^3 w_S^2 + 2\sigma^2 S (1 + 6\rho S w) w_S + \sigma^2 (1 + 6\rho S w) w = 0 \quad \text{in } \mathbb{R}_+ \times [0, \infty),$$

where $w = u_{SS}$.

Applying the transformations $w = \frac{v}{\rho S}$ and $x = \ln S$ to the reaction-advection-diffusion equation above and simplifying we get

$$v_t + \frac{\sigma^2}{2} (1 + 4v) v_{xx} + 2\sigma^2 v_x^2 + \frac{\sigma^2}{2} (1 + 4v) v_x = 0 \quad \text{in } \mathbb{R} \times [0, \infty). \tag{3.10}$$

If we let $v = \frac{V-1}{4}$ then (3.10) reduces to equation (3.1).

Hence, substituting $v = \frac{V-1}{4}$ into (3.8) gives the solution to (3.10) as

$$v(x, t) = \frac{1}{4}e^{\frac{x_0 - (x-ct)}{2}} + \frac{c}{\sigma^2} - \frac{1}{4}, \quad \sigma > 0, \quad t \geq 0. \tag{3.11}$$

Substituting $w = \frac{v}{\rho S}$ and $x = \ln S$ into (3.11) gives

$$u_{SS} = \frac{1}{\rho S} \left(\frac{1}{4} \sqrt{\frac{S_0}{S}} e^{\frac{ct}{2}} + \frac{c}{\sigma^2} - \frac{1}{4} \right), \quad \rho, c, S, S_0, \sigma > 0, \quad t \geq 0. \tag{3.12}$$

Since $\ln S_0 = x_0 \in \mathbb{R}$ then $S_0 > 0$.

We can rewrite equation (3.12) as

$$\rho S u_{SS} = \frac{1}{4} \sqrt{\frac{S_0}{S}} e^{\frac{ct}{2}} + \frac{c}{\sigma^2} - \frac{1}{4}. \tag{3.13}$$

By localizing boundary conditions we get $\lim_{S \rightarrow \pm\infty} u_{SS}(S, t) = 0$. Since

$$\lim_{S \rightarrow \pm\infty} u_{SS}(S, t) = \lim_{S \rightarrow \pm\infty} \frac{1}{\sqrt{S}} = 0,$$

equation (3.13) becomes $c = \frac{\sigma^2}{4} > 0$ after simplifying.

Plugging this value of c into (3.12) we obtain

$$u_{SS} = \frac{\sqrt{S_0}}{4\rho S^{3/2}} e^{\frac{\sigma^2 t}{8}}. \tag{3.14}$$

Integrate (3.14) twice with respect to S . After a little algebra we get

$$u(S, t) = aS - \frac{\sqrt{S_0}}{\rho} \left(\sqrt{S} e^{\frac{\sigma^2 t}{8}} + \frac{\sqrt{S_0}}{4} e^{\frac{\sigma^2 t}{4}} \right) \tag{3.15}$$

for $S, S_0, \sigma, \rho > 0, t \geq 0$ and $a \in \mathbb{R}$.

Substituting equation (3.15) into (2.5) and simplifying the results gives $a = 1$. Since $a = 1$ equation (3.15) reduces to (3.9). □

Remark 3.3. From equation (3.9), it is clear that $\frac{Su_S}{u}$ is always greater than one as in [1]. This shows that the option is always more volatile than the stock.

4. Conclusion

We have studied the hedging of derivatives in illiquid markets. A model where the implementation of a hedging strategy affects the price of the underlying asset has been considered. Assuming the solution of a forward wave, a classical solution for the nonlinear Black-Scholes equation was found. The solution obtained can be used for pricing a European *call* option at time $t \geq 0$. This solution supports the comments in [1] that the option is always more volatile than the stock.

In conclusion, further research needs to be done to solve the nonlinear Black-Scholes equation when $r > 0$. Future work will also involve finding solutions to arbitrary portfolio of options with non-single-signed gamma.

References

- [1] F. Black, M. Scholes, The pricing of options and corporate liabilities, *The Journal of Political Economy*, **81**, No. 3 (1973), 637-654.
- [2] U. Cetin, R. Jarrow, P. Protter, Liquidity risk and arbitrage pricing theory, *Finance Stoch.*, **8** (2004), 311-341.
- [3] J. Esekun, S. Onyango, N.O. Ongati, Analytic solution of a nonlinear Black-Scholes partial differential equation, *International Journal of Pure and Applied Mathematics*, **61**, No. 2 (2010), 219-230.
- [4] J.D. Logan, *An Introduction to Nonlinear Partial Differential Equations*, Second Edition, John Wiley and Sons, Inc., Hoboken, New Jersey, USA (2008).

