

A CHARACTERIZATION OF $PSU(19, q)$

A. Iranmanesh[§]

Department of Mathematics
Tarbiat Modarres University
P.O. Box 14115-137, Tehran, IRAN
e-mail: iranmana@modares.ac.ir

and

Institute for Studies in Theoretical Physics and Mathematics
Tehran, IRAN

Abstract: Let G be a finite group. The order of G is the product of coprime positive integers which is called the order components of G . It was proved that some non-abelian simple groups are uniquely determined by their order components. As the main result of this paper, we show that the simple groups $PSU(19, q)$ are also uniquely determined by their order components. As corollaries of this result, the validity of a conjecture of J.G. Thompson on $PSU(19, q)$ is obtained.

AMS Subject Classification: 20D06, 20D60

Key Words: simple group, prime graph, order component, finite group

1. Introduction

For an integer n , let $\pi(n)$ be the set of prime divisors of n . If G is a finite group then $\pi(G)$ is defined to be $\pi(|G|)$. The prime graph $\Gamma(G)$ of a group G is a graph whose vertex set is $\pi(G)$, and two distinct primes p and q are linked by an

Received: July 6, 2004

© 2004, Academic Publications Ltd.

[§]Correspondence address: Department of Mathematics, Tarbiat Modarres University, P.O. Box 14115-137, Tehran, IRAN

edge if and only if G contains an element of order pq . Let π_i , $i = 1, 2, \dots, t(G)$ be the connected components of $\Gamma(G)$. For $|G|$ even, π_1 will be the connected component containing 2. Then $|G|$ can be expressed as a product of some positive integers m_i , $i = 1, 2, \dots, t(G)$ with $\pi(m_i) = \pi_i$. The integers m_i 's are called the order components of G . The set of order components of G will be denoted by $OC(G)$. If the order of G is even, we will assume that m_1 is the even order component and $m_2, \dots, m_{t(G)}$ will be the odd order components of G . The order components of non-Abelian simple groups having at least three prime graph components are obtained by G.Y. Chen [5, Tables 1, 2, 3]. The order components of non-Abelian simple groups with two order components can be obtained according to [14, 15] (See [8, 9]). The following groups are uniquely determined by their order components: Sporadic simple groups, [2]; $PSL_2(q)$, [5]; $PSL(3, q)$, [8, 9]; $PSL(5, q)$, [7]; $C_2(q)$, where $q > 5$ [10]; $PSU(3, q)$ for $q > 5$, [13]; $PSU(5, q)$, [11]; $PSU(11, q)$, [12].

In this paper, we prove that $PSU(19, q)$ are also uniquely determined by their order components, that is we have the following theorem.

Main Theorem. *Let G be a finite group and $M = PSU(19, q)$. Then $OC(G) = OC(M)$ if and only if $G \cong M$.*

2. Preliminary Results

In order to prove the main theorem, first we bring some lemmas.

Definition 2.1. (see [6]) A finite group G is called a 2-Frobenius group if it has a normal series $1 < H < K < G$, where K and G/H are Frobenius groups with kernels H and K/H , respectively.

Lemma 2.2. (see [15], Theorem A) *If G is a finite group with its prime graph having more than one component, then G is one of the following groups:*

- (a) a Frobenius or 2-Frobenius group;
- (b) a simple group;
- (c) an extension of a π_1 -group by a simple group;
- (d) an extension of a simple group by a π_1 -solvable group;
- (e) an extension of a π_1 -group by a simple group by a π_1 -group.

Lemma 2.3. (see [15], Lemma 3) *If G is a finite group with more than one prime graph component and has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is simple, then H is a nilpotent group.*

The next lemma follows from Theorem 2 in [1].

Lemma 2.4. *Let G be a Frobenius group of even order and let H , K be Frobenius complement and Frobenius kernel of G , respectively. Then $t(G) = 2$,*

and the prime graph components of G are $\pi(H)$, $\pi(K)$ and, G has one of the following structures:

(a) $2 \in \pi(K)$ and all Sylow subgroups of H are cyclic.

(b) $2 \in \pi(H)$, K is an Abelian group, H is a solvable group, the Sylow subgroups of odd order of H are cyclic groups and the 2-Sylow subgroups of H are cyclic or generalized quaternion groups.

(c) $2 \in \pi(H)$, K is an Abelian group and there exists $H_0 \leq H$ such that $|H : H_0| \leq 2$, $H_0 = Z \times SL(2, 5)$, $(|Z|, 2.3.5) = 1$ and the Sylow subgroups of Z are cyclic.

The next lemma follows from Theorem 2 in [1] and Lemma 2.3.

Lemma 2.5. *Let G be a 2-Frobenius group of even order. Then $t(G) = 2$ and G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that:*

(a) $\pi_1 = \pi(G/K) \cup \pi(H)$ and $\pi(K/H) = \pi_2$;

(b) G/K and K/H are cyclic, $|G/K|$ divides $|\text{Aut}(K/H)|$, $(|G/K|, |K/H|) = 1$ and $|G/K| < |K/H|$;

(c) H is nilpotent and G is a solvable group.

Lemma 2.6. (see [4], Lemma 8) *Let G be a finite group with $t(G) \geq 2$ and let N be a normal subgroup of G . If N is a π_i -group for some prime graph component of G and m_1, m_2, \dots, m_r are some order components of G but not a π_i -number, then $m_1 m_2 \dots m_r$ is a divisor of $|N| - 1$.*

Lemma 2.7. (see [3], Lemma 1.4) *Suppose G and M are two finite groups satisfying $t(M) \geq 2$, $N(G) = N(M)$, where $N(G) = \{n \mid G \text{ has a conjugacy class of size } n\}$, and $Z(G) = 1$. Then $|G| = |M|$.*

The next lemma follows from Lemma 1.5 in [3].

Lemma 2.8. *Let G_1 and G_2 be finite groups satisfying $|G_1| = |G_2|$ and $N(G_1) = N(G_2)$. Then $t(G_1) = t(G_2)$ and $OC(G_1) = OC(G_2)$.*

Lemma 2.9. *Let G be a finite group and let M be a non-Abelian simple group with $t(M) = 2$ satisfying $OC(G) = OC(M)$. Let $|M| = m_1 m_2$, $OC(M) = \{m_1, m_2\}$, and $\pi(m_i) = \pi_i$ for $i=1$ or 2 . Then $|G| = m_1 m_2$ and one of the following holds:*

(a) G is a Frobenius or 2-Frobenius group;

(b) G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that G/K is a π_1 -group, H is a nilpotent π_1 -group, and K/H is a non-Abelian simple group. Moreover $OC(K/H) = \{m'_1, m'_2, \dots, m'_s, m_2\}$, $|K/H| = m'_1 m'_2 \dots m'_s m_2$ and $m'_1 m'_2 \dots m'_s \mid m_1$ where $\pi(m'_j) = \pi'_j$, $1 \leq j \leq s$. Also we have $|G/K| \mid |\text{Out}(K/H)|$.

Proof. The first part of the lemma follows from the above lemmas. Since $t(G) \geq 2$, we have $t(G/H) \geq 2$. Otherwise $t(G/H) = 1$, so that $t(G) = 1$. Moreover, we have $Z(G/H) = 1$. For any $xH \in G/H$ and $xH \notin K/H$, xH

GCD	$q + 1$	$q - 1$	$q^2 + 1$	$q^4 + 1$	$q^8 + 1$	c_1	c_2	c_3	c_4
$q + 1$	$q + 1$	1 or 2	1 or 2	1 or 2	1 or 2	1 or 11	1 or 7	1	1 or 5
$q - 1$	1 or 2	$q - 1$	1 or 2	1 or 2	1 or 2	1	1	1	1
$q^2 + 1$	1 or 2	1 or 2	$q^2 + 1$	1 or 2	1 or 2	1	1	1	1
$q^4 + 1$	1 or 2	1 or 2	1 or 2	$q^4 + 1$	1 or 2	1	1	1	1
$q^8 + 1$	1 or 2	1 or 2	1 or 2	1 or 2	$q^8 + 1$	1	1	1	1

Table 1:

induces an automorphism of K/H and this automorphism is trivial if and only if $xH \in Z(G/H)$. Therefore, $G/K \leq \text{Out}(K/H)$ and since $Z(G/H) = 1$, it follows that $|G/K| \mid |\text{Out}(K/H)|$. \square

Lemma 2.10. *Let $M = PSU(19, q)$. Suppose $D(q) = \frac{q^{19}+1}{k(q+1)}$, where $k = (19, q + 1)$. Then:*

- (a) *If $p \in \pi(M)$, then $|S_p| \leq q^{171}$, where $S_p \in \text{Syl}_p(M)$;*
- (b) *If $p \in \pi_1(M)$ and $p^\alpha \mid |M|$, then $p^\alpha - 1 \equiv 0 \pmod{D(q)}$ if and only if $p^\alpha = q^{38}, q^{76}, q^{114}$ or q^{152} ;*
- (c) *If $p \in \pi_1(M)$ and $p^\alpha \mid |M|$, then $p^\alpha + 1 \equiv 0 \pmod{D(q)}$ if and only if $p^\alpha = q^{19}, q^{57}, q^{95}, q^{133}$ or q^{171} .*

Proof. (a) From Table 1 in [8] we have $|M| = q^{171}(q + 1)^{18}(q - 1)^9(q^2 - q + 1)^6(q^2 + 1)^4(q^4 - q^3 + q^2 - q + 1)^3(q^2 + q + 1)^3 \times (1 - q + q^2 - q^3 + q^4 - q^5 + q^6)^2(q^4 + 1)^2(q^6 - q^3 + 1)^2(q^4 + q^3 + q^2 + q + 1) \times (1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10})(q^4 - q^2 + 1)(q^6 + q^3 + 1) \times (1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10} - q^{11} + q^{12}) \times (q^6 + q^5 + q^4 + q^3 + q^2 + q + 1)(1 + q - q^3 - q^4 - q^5 + q^7 + q^8)(q^8 + 1) \times (1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10} - q^{11} + q^{12} - q^{13} + q^{14} - q^{15} + q^{16}) \times \frac{(q^{19}+1)}{k(q+1)}$.

For convenience, let $c_1 := 1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10}$, $c_2 := 1 - q + q^2 - q^3 + q^4 - q^5 + q^6$, $c_3 := q^4 - q^2 + 1$, $c_4 := q^4 - q^3 + q^2 - q + 1$, $c_5 := q^4 + q^3 + q^2 + q + 1$, $c_6 := q^6 - q^3 + 1$, $c_7 := q^2 + q + 1$, $c_8 := q^2 - q + 1$, $c_9 := 1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10} - q^{11} + q^{12}$, $c_{10} := 1 + q - q^3 - q^4 - q^5 + q^7 + q^8$, $c_{11} := q^6 + q^5 + q^4 + q^3 + q^2 + q + 1$, $c_{12} := q^6 + q^3 + 1$, and $c_{13} := 1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10} - q^{11} + q^{12} - q^{13} + q^{14} - q^{15} + q^{16}$.

Now by easy calculations we can compute the greatest common divisors of every pair of

$$\{ q, q - 1, q + 1, q^2 + 1, q^4 + 1, q^8 + 1, c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8, c_9, c_{10}, c_{11}, c_{12}, c_{13} \}.$$

Obviously q is coprime with respect to another factors of $|M|$. In Table 1 and Table 2 we present some of these results.

GCD	c_5	c_6	c_7	c_8	c_9	c_{10}	c_{11}	c_{12}	c_{13}
$q + 1$	1	1 or 3	1	1 or 3	1 or 13	1	1	1	1 or 17
$q - 1$	1 or 5	1	1 or 3	1	1	1	1 or 7	1 or 3	1
$q^2 + 1$	1	1	1	1	1	1	1	1	1
$q^4 + 1$	1	1	1	1	1	1	1	1	1
$q^8 + 1$	1	1	1	1	1	1	1	1	1

Table 2:

By easy calculations we determine the greatest common divisors of any two factors of $|M|$.

Now let $p^\alpha \mid |M|$ and $p \in \pi_1$. As we mentioned above we can claim that one of the following occurs: p^α is a divisor of q^{171} , $2^{16} \times 3^8 \times 5^3 \times 7^2 \times 11 \times 13 \times 17 \times (q + 1)^{18}$, $2^{25} \times 3^4 \times 5 \times 7 \times (q - 1)^9$, $3^{20}(q^2 - q + 1)^6$, $2^{30}(q^2 + 1)^4$, $3^{10}(q^2 + q + 1)^3$, $2^{30}(q^4 + 1)^2$, $5^{18}(q^4 - q^3 + q^2 - q + 1)^3$, $5^9(q^4 + q^3 + q^2 + q + 1)$, $(q^4 - q^2 + 1)$, $3^{24}(q^6 - q^3 + 1)^2$, $3^{12}(q^6 + q^3 + 1)$, $7^9(q^6 + q^5 + q^4 + q^3 + q^2 + q + 1)$, $7^{18}(1 - q + q^2 - q^3 + q^4 - q^5 + q^6)^2$, $(1 + q - q^3 - q^4 - q^5 + q^7 + q^8)$, $2^{33}(q^8 + 1)$, $11^{18}(1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10})$, $13^{18}(1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10} - q^{11} + q^{12})$, $17^{18}(1 - q + q^2 - q^3 + q^4 - q^5 + q^6 - q^7 + q^8 - q^9 + q^{10} - q^{11} + q^{12} - q^{13} + q^{14} - q^{15} + q^{16})$.

Therefore the order of every Sylow subgroup of G is less than or equal to q^{171} , and hence (a) follows.

(b) Let there exists $p \in \pi_1(M)$, $p^\alpha \mid |M|$ and $p^\alpha - 1 \equiv 0 \pmod{D(q)}$. It is obvious that $p^\alpha > D(q)$. Similar to the proof of (a) we must consider different cases. For $q \leq 27$ numerical calculations show that there is no p^α such that (b) holds. Hence we can let $q > 27$. But it is straightforward to see that if $q \geq 29$, then $D(q) > (q + 1)^{18}/38$.

We consider the following cases:

(1) If $p^\alpha \mid 2^{16} \times 3^8 \times 5^3 \times 7^2 \times 11 \times 13 \times 17 \times (q + 1)^{18}$, then we consider the following subcases:

(1.1) Let $p \neq 2, 3, 5, 7, 11, 13, 17$ and $p^\alpha \mid (q + 1)^{18}$ and $p^\alpha - 1 \equiv 0 \pmod{D(q)}$, then $p^\alpha - 1 = sD(q)$ for some $s > 0$. But $(q + 1)^{18}/38 < D(q)$, which implies that $p^\alpha = (q + 1)^{18}/t$, where $st \leq 38$. Now, numerical calculations show that these equations have no a solution in \mathbb{Z} and hence, there can not exist any p and α such that the above relations holds.

(1.2) If $p = 2$, then $2^\alpha \mid 2^{16}(q + 1)^{18}$. Hence $2^{16}(q + 1)^{18}/t - 1 = sD(q)$, where $st \leq 2^{17} \times 19$, since $2^{16}(q + 1)^{18} < 2^{17} \times 19D(q)$ for $q > 27$. By expanding the above equation we can get a diophantine equation and by solving this equation we see that there exist no α such that (b) holds.

(1.3) If $p = 3, 5, 7, 11, 13$ or 17 , then $3^\alpha | 3^8(q+1)^{18}$, $5^\alpha | 5^3(q+1)^{18}$, $7^\alpha | 7^2(q+1)^{18}$, $11^\alpha | 11(q+1)^{18}$, $13^\alpha | 13(q+1)^{18}$, $17^\alpha | 17(q+1)^{18}$, respectively. We get a contradiction similar to (1.2).

(2) If $p^\alpha | 2^{25} \times 3^4 \times 5 \times 7 \times (q-1)^9$, then p^α divides $2^{25}(q-1)^9$, $3^4(q-1)^9$, $5(q-1)^9$ or $7(q-1)^9$. But in each case $p^\alpha < D(q)$ which implies that $p^\alpha - 1 \not\equiv 0 \pmod{D(q)}$.

(3) If p^α is a divisor of $3^{20}(q^2-q+1)^6$, $3^{10}(q^2+q+1)^3$, $2^{30}(q^2+1)^4$, $2^{33}(q^4+1)^2$, $5^{18}(q^4-q^3+q^2-q+1)^3$, $5^9(q^4+q^3+q^2+q+1)$, (q^4-q^2+1) , $7^{18}(1-q+q^2-q^3+q^4-q^5+q^6)^2$, $3^{24}(q^6-q^3+1)^2$, $3^{12}(q^6-q^3+1)$, $7^9(q^6+q^5+q^4+q^3+q^2+q+1)$, $(1+q-q^3-q^4-q^5+q^7+q^8)$, $2^{33}(q^8+1)$, $11^{18}(1-q+q^2-q^3+q^4-q^5+q^6-q^7+q^8-q^9+q^{10})$, $13^{18}(1-q+q^2-q^3+q^4-q^5+q^6-q^7+q^8-q^9+q^{10}-q^{11}+q^{12})$, $17^{18}(1-q+q^2-q^3+q^4-q^5+q^6-q^7+q^8-q^9+q^{10}-q^{11}+q^{12}-q^{13}+q^{14}-q^{15}+q^{16})$, then in each case $p^\alpha < D(q)$ which implies that $p^\alpha - 1 \not\equiv 0 \pmod{D(q)}$.

(4) At last, let $p^\alpha | q^{171}$. Then we consider two subcases, namely $k = 1$ and $k = 19$. Since the proofs are similar we state only one of them, namely $k = 1$ and the other case is similar.

We know that $q = p^n$ for some $n > 0$.

First, we prove that if $p^\beta | q^{19}$ and $p^\beta + 1 \equiv 0 \pmod{D(q)}$, then $p^\beta = q^{19}$. We have

$$p^\beta + 1 = s \cdot D(q) = s \cdot \frac{q^{19} + 1}{q + 1} = s(q^{18} - q^{17} + q^{16} - \dots + q^2 - q + 1),$$

which implies that $1 \leq s \leq q + 1$, since $p^\beta \leq q^{19}$. Also since $q | p^\beta$ we have $q | s - 1$ which implies that $q \leq s - 1$. Therefore $q = s$ and hence $p^\beta = q^{19}$.

Now we prove that if $p^\alpha | q^{38}$ and $p^\alpha - 1 \equiv 0 \pmod{D(q)}$, then $p^\alpha = q^{38}$. Now we consider two cases. First let $p^\alpha \leq q^{19}$ and $p^\alpha - 1 \equiv 0 \pmod{D(q)}$. In this case $p^\alpha - 1 = s \cdot D(q)$, where $s < q + 1$. Since $q | p^\alpha$ we have $q | s + 1$ and hence $q \leq s + 1$. Therefore $s = q$ or $s = q - 1$. But easy calculation shows that in each case $p^\alpha - 1 \neq s \cdot D(q)$, which is a contradiction. Therefore $p^\alpha > q^{19}$ and hence $p^\alpha = q^{19} p^m$ for some $m > 0$. Thus we have

$$p^\alpha - 1 = q^{19} p^m - 1 = p^m (q^{19} + 1) - p^m - 1.$$

Therefore $D(q) | p^m + 1$ which implies that $p^m = q^{19}$, by the above statement and hence $p^\alpha = q^{38}$.

If $p^\alpha > q^{38}$ and $p^\alpha | q^{171}$, then by a similar method we conclude that $p^\alpha = q^{76}$, q^{114} or q^{152} .

(c) Similar to part (b) we conclude that p^α must be equal to q^{19} , q^{57} , q^{95} , q^{133} or q^{171} and the proof is completed. \square

Remark. In the sequel of this paper and specially in the proof of the main theorem, for convenience we suppose that $X = \{q^{38}, q^{76}, q^{114}, q^{152}\}$ and $Y = \{q^{19}, q^{57}, q^{95}, q^{133}, q^{171}\}$. Therefore if $p \in \pi_1(M)$ and $p^\alpha \mid |M|$, then $p^\alpha - 1 \equiv 0 \pmod{D(q)}$ if and only if $p^\alpha \in X$, and $p^\alpha + 1 \equiv 0 \pmod{D(q)}$ if and only if $p^\alpha \in Y$.

Lemma 2.11. *Let G be a finite group and $M = PSU(19, q)$ and $OC(G) = OC(M)$. Then G is neither a Frobenius group nor a 2-Frobenius group.*

Proof. We will use some results about Frobenius groups. For example we know that if G is a Frobenius group, by Lemma 2.4, $OC(G) = \{|H|, |K|\}$, where K and H are the Frobenius kernel and the Frobenius complement of G , respectively. Also we know that $|H| \mid (|K| - 1)$, and hence $|H| < |K|$. So $|H| = \frac{q^{19}+1}{(q+1)(19, q+1)}$, $|K| = |G|/|H|$. There exists a prime p such that $p^\alpha \mid 7(q-1)^9$. If P is a p -Sylow subgroup of K , then since K is nilpotent, $P \triangleleft G$ and hence $D(q) \mid |P| - 1$ by Lemma 2.6, which implies that $p^\alpha \in Y$ by Lemma 2.10 (b). But obviously $7(q-1)^9 < q^{38}$ which is a contradiction. Therefore, G is not a Frobenius group.

Let G be a 2-Frobenius group. By Lemma 2.5, there is a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that $|K/H| = \frac{q^{19}+1}{(q+1)(19, q+1)} < 2^{16}(q+1)^{18}$ and $|G/K| < |K/H|$. Thus there exists a prime p such that $p \mid 2^{16}(q+1)^{18}$ and $p \mid |H|$. If P is a p -Sylow subgroup of H , since H is nilpotent, P must be a normal subgroup of K with $P \subseteq H$ and $|K| = \frac{q^{19}+1}{k(q+1)}|H|$. Therefore, $\frac{q^{19}+1}{k(q+1)} \mid (|P| - 1)$, by Lemma 2.6 and hence $q^{38} \mid |P|$, which is impossible since $|P| \leq 2^{16}(q+1)^{18}$. Therefore, G is not a 2-Frobenius group. □

Lemma 2.12. *Let G be a finite group. If the order components of G are the same as those of $M = PSU(19, q)$, then G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is a simple group. Moreover, the odd order component of M is equal to some of those of K/H , and in particular, $t(K/H) \geq 2$.*

Proof. The first part of the lemma follows from the above lemmas since the prime graph of M has two components. For primes p and q , if K/H has an element of order pq , then G has one. Hence, by the definition of prime graph component, the odd order component of G must be an odd order component of K/H . □

3. Proof of Main Theorem

By Lemma 2.12, G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups, K/H is a non-Abelian simple group, $t(K/H) \geq 2$ and the odd order component of M is an odd order component of K/H .

We now proceed the proof in the following steps:

Step 1. If $K/H \cong A_n$, where $n = p, p + 1, p + 2$ and $p \geq 5$ is a prime number, then we have two cases:

Case 1. $k = 1$. In this case, p or $p - 2$ are equal to $\frac{q^{19}+1}{q+1}$. If $p = \frac{q^{19}+1}{q+1}$, then $p - 1 = q(q - 1)(q^2 - q + 1)(q^2 + q + 1)(q^6 - q^3 + 1)(q^6 + q^3 + 1)$ and

$$p - 2 = q^{18} - q^{17} + q^{16} - q^{15} + q^{14} - q^{13} + q^{12} - q^{11} + q^{10} - q^9 + q^8 - q^7 + q^6 - q^5 + q^4 - q^3 + q^2 - q - 1. \quad (1)$$

But easy calculation shows that $(p - 2, |G|) \mid 31 \times 17$. But for $q > 1$, $D(q) > 31 \times 17$ and it is a contradiction.

If $p - 2 = \frac{q^{19}+1}{q+1}$, then we proceed similarly for $p - 4$ since $p > 5$.

Case 2. $k = 19$. Then p or $p - 2$ is equal to $\frac{q^{19}+1}{19(q+1)}$ and $p - 2$ or $p - 4$ must be equal to $\frac{q^{18}-q^{17}+q^{16}-q^{15}+\dots-q-37}{19}$, respectively. Now we proceed similarly to the last case and get a contradiction.

Step 2. If K/H is a sporadic simple group, then $D(q)$ must be equal to 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 59, 67, 71, which has no solution, since $D(2) = 174763$.

Therefore K/H is a simple group of Lie type.

Step 3. If $K/H \cong E_6(q')$ or ${}^2E_6(q')$, then we get a contradiction.

Step 4. If $K/H \cong A_r(q')$, then we distinguish the following 6 cases:

4.1. $K/H \cong A_{p'-1}(q')$, where $(p', q') \neq (3, 2), (3, 4)$.

In fact we have $D(q) = (q'^{p'} - 1)/((q' - 1)(p', q' - 1))$ and hence $q'^{p'} - 1 \equiv 0 \pmod{D(q)}$. Now by using Lemma 2.10(b) we have $q'^{p'} \in X$, which implies that $q'^{p'} = q^{38}, q^{76}, q^{114}$ or q^{152} . If $p' \geq 11$, then $q'^{\frac{p'(p'-1)}{2}} > q^{171}$, which is impossible by Lemma 2.11(a). Therefore we must check cases $p' = 3, 5$ and 7. If $p = 3$ and $q'^3 = q^{38}$, then

$$(q^{19} - 1)(q + 1)(19, q + 1) = (q' - 1)(3, q' - 1), \quad q'^3 = q^{38}.$$

But these equations have no common solution in \mathbb{Z} , and hence this case is also impossible. If $p' = 3$ and $q'^3 = q^{76}, q^{114}$ or q^{152} ; or if $p' = 5, 7$, then we get a contradiction similarly.

4.2. $K/H \cong A_{p'}(q')$, where $(q' - 1)|(p' + 1)$. Then $q'^{p'} \in X$. But for $p' > 7$ we have $q'^{\frac{p'(p'+1)}{2}} > q^{171}$, which is impossible. If $p' = 3, 5$ or 7 , then $q' - 1 < 8$, since $q' - 1 \mid p' + 1$. Now easily we can get a contradiction.

4.3. $K/H \cong A_1(q')$, where $4|(q' + 1)$.

The odd order components of K/H are q' and $(q' - 1)/2$.

If $D(q) = \frac{q'-1}{2}$, then $q' \in X$, which implies that $q' = q^{38}, q^{76}, q^{114}$ or q^{152} . If $q' = q^{38}$, then we have $(q19+1)/k(q+1) = \frac{q^{38}-1}{2}$. Therefore, $2 = k(q+1)(q^{19}-1)$ and it is impossible.

If $D(q) = q'$, then by similar method in Step 1, we get a contradiction.

4.4. $K/H \cong A_1(q')$, where $4|(q' - 1)$.

The odd order components of K/H are q' and $(q' + 1)/2$.

If $D(q) = \frac{q'+1}{2}$, then $q' \in Y$, which implies that $q' = q^{19}, q^{57}, q^{95}, q^{133}$ or q^{171} . By similar method in (4.3), we get a contradiction

If $D(q) = q'$ then we proceed similarly to above case and get a contradiction.

4.5. $K/H \cong A_1(q')$, where $4 \mid q'$.

The odd order components of K/H are $q' + 1$ and $q' - 1$. By similar method in last case, we get a contradiction.

4.6. $K/H \cong A_2(2)$ or $K/H \cong A_2(4)$. Then $D(q)$ must be equal to 3, 5, 7, 9 which is impossible.

Step 5. If $K/H \cong B_r(q')$, or $C_r(q')$, or $D_r(q')$, or $F_4(q')$, or ${}^3D_4(q')$, or $E_8(q')$, or ${}^2G_2(q')$, by a similar method we get contradictions.

Step 6. If $K/H \cong E_7(2), E_7(3), {}^2E_6(2),$ or ${}^2F_4(2)'$, then $D(q)$ must be equal to 13, 17, 19, 73, 127, 757, 1093 which is impossible.

Step 7. If $K/H \cong G_2(q')$, then we consider 3 cases:

7.1. $K/H \cong G_2(q')$, where $2 < q' \equiv 1 \pmod{3}$. Then $D(q) = q'^2 - q' + 1$ and hence $q'^3 \in Y$, which implies that $q'^3 = q^{19}, q^{57}, q^{95}, q^{133}$ or q^{171} . If $q'^3 = q^{19}$, then

$$\frac{q'^3 + 1}{q' + 1} = \frac{q^{19} + 1}{k(q + 1)}.$$

Obviously $k = 1$ implies that $q = q'$ which is impossible. If $k = 19$, then $q^{19} = (19q + 18)^3$, which has no solution in \mathbb{Z} .

7.2. $K/H \cong G_2(q')$, where $2 < q' \equiv -1 \pmod{3}$. Then $D(q) = q'^2 + q' + 1$ and hence $q'^3 \in X$. Now we can proceed similar to 7.1 and get contradiction.

7.3. $K/H \cong G_2(q')$, where $3 \mid q'$. Then $q'^2 \pm q' + 1 = D(q)$. This is similar to case 7.1 and case 7.2.

Step 8. If $K/H \cong {}^2F_4(q')$, where $q' = 2^{2r+1} > 2$, or ${}^2B_2(q')$, where $q' = 2^{2t+1} > 2$ we can get a contradiction by a similar method in above step.

Step 9. If $K/H \cong {}^2D_r(q')$, then we consider 6 cases:

- 9.1. $K/H \cong {}^2D_r(q')$, where $r = 2^t \geq 4$.
- 9.2. $K/H \cong {}^2D_r(2)$, where $r = 2^t + 1 \geq 5$.
- 9.3. $K/H \cong {}^2D_p(3)$, where $5 \leq p \neq 2^r + 1$.
- 9.4. $K/H \cong {}^2D_r(3)$, where $r = 2^t + 1 \neq p$, $t \geq 2$.
- 9.5. $K/H \cong {}^2D_p(3)$, where $p = 2^t + 1$, $t \geq 2$.
- 9.6. $K/H \cong {}^2D_{p+1}(2)$, where $p = 2^r - 1$, $r \geq 2$.

In all of above cases, we get a contradiction.

Step 10. If $K/H \cong {}^2A_r(q')$, then we consider 3 cases:

10.1. $K/H \cong {}^2A_3(2)$, ${}^2A_3(3)$ or ${}^2A_5(2)$. Then $D(q)$ must be equal to 5, 7, 11 which is impossible.

10.2. $K/H \cong {}^2A_{p'}(q')$, where $(q' + 1)|(p' + 1)$ and $(p', q') \neq (3, 3), (5, 2)$. Then $D(q) = q^{p'} + 1/q' + 1$ and hence $q^{p'} \in Y$ which implies that $q^{p'} = q^{19}, q^{57}, q^{95}, q^{133}$ or q^{171} . If $\frac{p'+1}{2} > 9$, then $q^{p(p+1)/2} > q^{171}$ which is a contradiction, by Lemma 2.10(a). Therefore $p' = 3, 5, 7, 11, 13, 17$. If $p' = 3$, then $q' = 3$, since $q' + 1|p' + 1$. But it is a contradiction, since $(p', q') \neq (3, 3)$.

If $p' = 5$, then $q' + 1|6$. Since $(p', q') \neq (5, 2)$, we have $q' = 5$. But $q^{19} = 5^5$, which is a contradiction.

Similarly we get a contradiction in other cases.

10.3. $K/H \cong {}^2A_{p'-1}(q')$. Then $q^{p'} = q^{19}, q^{57}, q^{95}, q^{133}$ or q^{171} . If $p' > 19$, then $q^{\frac{p'(p'-1)}{2}} > q^{171}$, which is impossible. Otherwise, if $p' = 3, 5, 7, 11, 13$ or 17 , then

$$(q' + 1)(p', q' + 1) = (q + 1)(19, q + 1), \quad q^{p'} = q^{19}.$$

But these equations have no common solution in \mathbb{Z} . If $p' = 19$, then $q = q'$. Thus $|G| = |PSU(19, q)| = |K/H| = |K|/|H|$ which implies that $|H| = 1$ and $|K| = |G| = |PSU(19, q)|$. Therefore, $K = PSU(19, q)$ and hence $G = PSU(19, q)$.

The proof of the main theorem is now completed.

Remark 3.1. It is a well-known conjecture of J.G. Thompson that if G is a finite group with $Z(G) = 1$ and M is a non-Abelian simple group satisfying $N(G) = N(M)$, then $G \cong M$.

We can give a positive answer to this conjecture for the groups under discussion.

Corollary 3.2. *Let G be a finite group with $Z(G) = 1$, $M = PSU(19, q)$ and $N(G) = N(M)$, then $G \cong M$.*

Proof. By Lemma 2.8 if G and M are two finite groups satisfying the conditions of Corollary 3.2, then $OC(G) = OC(M)$. So the main theorem implies this corollary. □

Acknowledgments

The author would like to thank the Institute for Studies in Theoretical Physics and Mathematics (IPM) Tehran, Iran for the financial support (No. 82200013).

References

- [1] G.Y. Chen, On Frobenius and 2-Frobenius group, *J. Southwest China Normal Univ.*, **20**, No. 5 (1995), 485-487.
- [2] G.Y. Chen, A new characterization of sporadic simple groups, *Algebra Colloq.*, **3**, No. 1 (1996), 49-58.
- [3] G.Y. Chen, On Thompson's conjecture, *J. Algebra*, **15** (1996), 184-193.
- [4] G.Y. Chen, Further reflections on Thompson's conjecture, *J. Algebra*, **218** (1999), 276-285.
- [5] G.Y. Chen, A new characterization of $PSL_2(q)$, *Southeast Asian Bulletin of Math*, **22** (1998), 257-263.
- [6] K.W. Gruenberg, K.W. Roggenkamp, Decomposition of the augmentation ideal and of the relation modules of a finite group, *Proc. London Math. Soc.*, **31** (1975), 146-166.
- [7] A. Iranmanesh, S.H. Alavi, A characterization of simple groups $PSL(5, q)$, *Bull. Austral. Math. Soc.*, **65** (2002), 211-222.
- [8] A. Iranmanesh, S.H. Alavi, B. Khosravi, A characterization of $PSL(3, q)$ where q is an odd prime power, *J. Pure and Applied Algebra*, **170**, No. 2-3 (2002), 243-254.
- [9] A. Iranmanesh, S.H. Alavi, B. Khosravi, A characterization of $PSL(3, q)$ for $q = 2^n$, *Acta Math. Sinica, English series*, **18**, No. 3 (2002), 463-472.
- [10] A. Iranmanesh, B. Khosravi, A characterization of $C_2(q)$ where $q > 5$, *Comment Math. Univ. Carolinae*, **43**, No. 1 (2002), 9-21.
- [11] A. Iranmanesh, B. Khosravi, A characterization of $PSU_5(q)$, *International Mathematical Journal*, **3**, No. 2 (2003), 129-141.
- [12] A. Iranmanesh, B. Khosravi, A characterization of $PSU_{11}(q)$, *Canadian Mathematical Bulletin*, To Appear.

- [13] A. Iranmanesh, B. Khosravi, S.H. Alavi, A characterization of $PSU(3, q)$ for $q > 5$, *Southeast Asian Bulletin Math*, **26**, No. 2 (2002), 33-44.
- [14] A.S. Kondrat'ev, Prime graph components of finite groups, *Math. USSR-sb.*, **67**, No. 1 (1990), 235-247.
- [15] J.S. Williams, Prime graph components of finite groups, *J. Algebra*, **69** (1981), 487-513.