

LINEAR SPACES OF MATRICES, SYMMETRIC  
MATRICES OR HERMITIAN MATRICES WITH  
A FIXED RANK OVER A FINITE FIELD

E. Ballico

Department of Mathematics  
University of Trento  
380 50 Povo (Trento) - Via Sommarive, 14, ITALY  
e-mail: ballico@science.unitn.it

**Abstract:** Here we raise several questions concerning linear spaces of matrices with fixed rank over  $\mathbb{F}_q$ .

**AMS Subject Classification:** 15A30, 12E20, 14N05

**Key Words:** linear spaces of matrices, rank, Hermitian matrices over a finite field

\*

Here we raise the following questions.

**Question 1.** Fix integers  $m \geq n \geq k > 0$ ,  $e \geq 1$ , and a prime power  $q$ . What is the maximal integer  $a(q, m, n, k)$ , (resp.  $b(q, m, n, k)$ ) such that there is an  $a(q, m, n, k)$ -dimensional (resp.  $b(q, m, n, k)$ -dimensional)  $\mathbb{F}_q$ -vector space of  $n \times m$  matrices over  $\mathbb{F}_q$  such that each non-zero element of it has rank  $k$  (resp.  $\leq k$ )? What is the maximal integer  $a(q, m, n, k, e)$ , (resp.  $b(q, m, n, k, e)$ ) such that there is an  $a(q, m, n, k, e)$ -dimensional (resp.  $b(q, m, n, k, e)$ -dimensional)  $\mathbb{F}_{q^e}$ -vector space of  $n \times m$  matrices with a basis defined over  $\mathbb{F}_q$  such that each non-zero element of it has rank  $k$  (resp.  $\leq k$ )? Say something about the linear subspaces with maximal dimension.

**Question 2.** Fix integers  $n \geq k > 0$ ,  $e \geq 1$ , and a prime power  $q$ . What is the maximal integer  $c(q, n, k)$ , (resp.  $d(q, n, k)$ ) such that there is a  $c(q, n, k)$ -dimensional (resp.  $d(q, n, k)$ -dimensional)  $\mathbb{F}_q$ -vector space of symmetric  $n \times n$

matrices over  $\mathbb{F}_q$  such that each non-zero element of it has rank  $k$  (resp.  $\leq k$ )? What is the maximal integer  $c(q, n, k, e)$ , (resp.  $d(q, m, n, k)$ ) such that there is an  $a(q, m, m, k, e)$ -dimensional  $\mathbb{F}_{q^e}$ -vector space of  $n \times m$  matrices with a basis defined over  $\mathbb{F}_q$  such that each non-zero element of it has rank  $k$  (resp.  $\leq k$ )? Say something about the linear subspaces with maximal dimension.

**Question 3.** Fix integers  $n \geq k > 0$  and a prime power  $q$ . What is the maximal integer  $e(q, n, k)$  (resp.  $f(q, n, k)$ ) such that there is an  $e(q, n, k)$ -dimensional (resp.  $f(q, n, k)$ -dimensional)  $\mathbb{F}_q$ -linear spaces of Hermitian matrices over  $\mathbb{F}_{q^2}$  such that each non-zero element of it has rank  $k$  (resp.  $\leq k$ )?

**Question 4.** Fix integers  $m \geq n \geq k > 0$  and a prime power  $q$  such that  $0 < t < a(q, m, n, k)$ . Show the existence of a  $t$ -dimensional  $\mathbb{F}_q$ -vector space of  $n \times m$  matrices over  $\mathbb{F}_q$  such that each non-zero element of it has rank  $k$ , but that it is not contained in a bigger linear space with the same property. Hopefully, prove the same guess for the other integers introduced in Questions 1, 2 and 3.

For background on Hermitian matrices over a finite field, see [6], Chapter 23; however, here we use the notation  $\mathbb{F}_q$  (resp.  $\mathbb{F}_{q^2}$ ) instead of  $\mathbb{F}_{\sqrt{q}}$  (resp.  $\mathbb{F}_q$ ). For similar problems for the real, complex and quaternionic division rings, see [1], [2], [3], [4], [5] and [7].

**Question 5.** Fix as the base field  $F$  either the real or the complex or the quaternionic division ring. Take the set-up of Questions 1 or 2 for  $e = 1$ . Is there a maximal dimension linear subspace defined over the rational field?

**Remark 1.** Fix as the base field  $F$  either the real or the complex or the quaternionic division ring. Take the case  $m = n$  and  $k = n - 1$ . The existence parts of the proofs in [1], [2] and [3] use an induction on  $n$  in which all the 3 division rings play simultaneously. In this way we get at least in terms of  $n$  an explicit upper bound for the degree of the extension  $L$  of  $\mathbb{Q}$  on which a maximal dimension linear subspace may be defined over  $L$ .

**Remark 2.** Take the case  $k = n$ . Let  $A, A_1, \dots, A_t$  be  $n \times m$  matrices over  $\mathbb{F}_q$ .  $\text{rank}(A) < n$  if and only if  $m - n + 1$   $n \times n$  submatrices of  $A$  have zero-determinant. Hence the set  $B(A_1, \dots, A_t) =$  of all  $(x_1, \dots, x_t) \in \mathbb{F}_{q^e}^{\oplus t}$  such that  $\text{rank}(x_1 A_1 + \dots + x_t A_t) < n$  is defined by  $m - n + 1$  homogeneous degree  $n$  equations with coefficients in  $\mathbb{F}_q$ . By Chevalley-Waring Theorem for systems of homogeneous equations we get  $a(q, m, n, n) \leq n(m - n + 1)$ . The same theorem gives an upper bound for the integer  $a(q, m, n, k)$  for any  $k < n$ . However, these upper bounds do not use the structure of the determinantal equations and hence they probably are very bad.

### Acknowledgements

The author was partially supported by MIUR and GNSAGA of INdAM (Italy).

### References

- [1] J. Adams, P. Lax, R. Phillips, On matrices whose real linear combinations are nonsingular, *Proc. Amer. Math. Soc.*, **16**, No. 2 (1965), 318-322.
- [2] Correction to “On matrices whose real linear combinations are nonsingular”, *Proc. Amer. Math. Soc.*, **17** (1966), 945-947.
- [3] Y. Au-Yeung, On matrices whose nontrivial real linear combinations are nonsingular, *Proc. Amer. Math. Soc.*, **29**, No. 1 (1971), 17-22.
- [4] A. Causin, G.P. Pirola, A note on spaces of symmetric matrices, *ArXiv: math.AT/0702270*.
- [5] D. Falikman, S. Friedland, R. Loewry, On spaces of matrices containing a nonzero matrix of bounded rank, *Pacific J. Math.*, **207**, No. 1 (2002), 157-176.
- [6] J.W.P. Hirschfeld, J.A. Thas, *General Galois Geometries*, Clarendon Press, Oxford (1991).
- [7] E.G. Rees, Linear spaces of real matrices of given rank, *Contemp. Math.*, **188** (1995), 219-229.
- [8] C. Small, *Arithmetic of Finite Fields*, Marcel Dekker, Inc., New York (1991).

