

OSCILLATORY PROPERTIES OF THE SOLUTIONS OF
A FIRST ORDER NEUTRAL NONCONSTANT DELAY
IMPULSIVE DIFFERENTIAL EQUATIONS WITH
VARIABLE COEFFICIENTS

M.B. Dimitrova^{1 §}, V.I. Donev²

^{1,2}Department of Mathematics
Technical University of Sliven
8800, Sliven, BULGARIA

Abstract: In this paper we consider first order neutral impulsive differential equation with variable coefficients. The asymptotic behavior of a non-oscillatory solution for such equations is investigating and sufficient conditions for oscillation of all the solutions are found.

AMS Subject Classification: 34A37, 34K11, 34K40, 34K45

Key Words: oscillation, impulsive effect, neutral differential equation

1. Introduction

The Neutral Impulsive Differential Equations (NIDE) are part of the Impulsive Differential Equations with Deviating Arguments (IDEDA). Among the numerous publications concerning the oscillation properties of IDEDA - with delayed or advanced arguments, we choose to refer to [1], [2], [8], [9], [10], [14] and [15]. NIDE are characterized with neutral argument in which the highest-order derivative of the unknown function appears in the equation both with and without delay. Moreover, the unknown function in them, may have discontinuities of first kind at points, which we call jump points. Such equations can be used to model processes, that occur in the theory of optimal control, industrial robotics, biotechnologies, etc. Some results on the oscillation theory of this type of equations can be found in [3], [4] and [7].

Received: July 7, 2011

© 2011 Academic Publications, Ltd.

[§]Correspondence author

As concern the behavior of the solutions of the differential equations with deviating arguments, but without impulses, we choose to refer to [6], [11], [12] and [13].

The authors investigated neutral delay impulsive differential equations with constant coefficients and found there necessary and sufficient conditions for existence of eventually positive solutions in [3] and established oscillation criteria in [4], as well. In the present paper we study the asymptotic behavior of the eventually non-oscillatory solutions of (E_1) and obtain oscillation criteria when the coefficients are variable and the delays are nonconstant.

2. Preliminary notes

The object of investigation in the present work is the first order impulsive differential equation with variable coefficients and nonconstant neutral delay argument of the form

$$\frac{d}{dt}\{y(t) - c(t)y(h(t))\} + p(t)y(\sigma(t)) = 0, \quad t \neq \tau_k, \quad k \in N \tag{E_1}$$

$$\Delta\{y(\tau_k) - c_{\tau_k}y(h(\tau_k))\} + p_{\tau_k}y(\sigma(\tau_k)) = 0, \quad k \in N$$

as well as the corresponding to it inequalities

$$\frac{d}{dt}\{y(t) - c(t)y(h(t))\} + p(t)y(\sigma(t)) \leq 0, \quad t \neq \tau_k, \quad k \in N \tag{N_{1,\le}}$$

$$\Delta\{y(\tau_k) - c_{\tau_k}y(h(\tau_k))\} + p_{\tau_k}y(\sigma(\tau_k)) \leq 0, \quad k \in N$$

and

$$\frac{d}{dt}\{y(t) - c(t)y(h(t))\} + p(t)y(\sigma(t)) \geq 0, \quad t \neq \tau_k, \quad k \in N \tag{N_{1,\ge}}$$

$$\Delta\{y(\tau_k) - c_{\tau_k}y(h(\tau_k))\} + p_{\tau_k}y(\sigma(\tau_k)) \geq 0, \quad k \in N.$$

The points $\tau_k \in (0, +\infty)$, $k \in N$ are the moments of impulsive effect (let us call them jump points), where the unknown function reveals its discontinuities of first kind as jumps. In order to manifest these jumps of the unknown function $y(t)$, we use the notation

$$\Delta\{y(\tau_k) - c_{\tau_k}y(h(\tau_k))\} = \Delta y(\tau_k) - c_{\tau_k}\Delta y(h(\tau_k)), \quad \Delta y(\tau_k) = y(\tau_k+0) - y(\tau_k-0).$$

Denote by $P_\tau C(R, R)$ the set of all functions $u: R \rightarrow R$, which satisfy the following conditions:

- (i) u is piecewise continuous on $(\tau_k, \tau_{k+1}]$, $k \in N$,
- (ii) u is continuous from the left at the points τ_k , i.e.

$$u(\tau_k - 0) = \lim_{t \rightarrow \tau_k - 0} u(t) = u(\tau_k),$$

(iii) there exists a sequence of reals $\{u(\tau_k + 0)\}_{k=1}^\infty$, such that

$$u(\tau_k + 0) = \lim_{t \rightarrow \tau_k + 0} u(t),$$

(iv) u may have discontinuities of first kind at the jump points $\tau_k, k \in N$, that we

qualify as down-jumps when $\Delta u(\tau_k) < 0$, or as up-jumps when $\Delta u(\tau_k) > 0, k \in N$.

Introduce the following hypotheses, where $R^+ = (0, +\infty)$ and $R_0^+ = [0, +\infty)$:

(H₁) $0 < \tau_1 < \tau_2 < \dots < \tau_k < \dots, \lim_{k \rightarrow +\infty} \tau_k = +\infty, \max\{\tau_{k+1} - \tau_k\} < +\infty, k \in N$;

(H₂) $h, \sigma \in C^1(R^+, R^+), h'(t) > 0, \sigma'(t) > 0, \sigma(t) < t, h(t) < t$
and $\lim_{t \rightarrow +\infty} h(t) = +\infty, \lim_{t \rightarrow +\infty} \sigma(t) = +\infty$;

(H₃) $c \in C(R, (0, c_0]), c_0 \in (0, 1), c_{\tau_k} = c(\tau_k)$;

(H₄) $p \in P_\tau C(R, R), p(t)$ is not identically zero on any positive half-line,
 $\sum_{k=1}^{+\infty} p_{\tau_k}^2 \neq 0$;

(H₅) $p \in P_\tau C(R, R^+), p_{\tau_k} \in R_0^+, k \in N, \int_0^{+\infty} p(s)ds + \sum_{k=1}^{+\infty} p_{\tau_k} = +\infty$.

Let $\rho(t) = \min_{t \in R^+} \{\sigma(t), h(t)\}$. We say that a real valued function $y(t)$ is a *solution* of the equation (E_1) , if there exists a number $T_0 \in R$ such that $y \in P_\tau C([\rho(T_0), +\infty), R)$, the function $z(t) = y(t) - c(t)y(h(t))$ is continuously differentiable for $t \geq T_0, t \neq \tau_k, k \in N$ and $y(t)$ satisfies (E_1) for all $t \geq T_0$.

Without further mentioning we will assume throughout this paper, that every solution $y(t)$ of equation (E_1) that is under consideration here, is continuable to the right and is nontrivial. That is, $y(t)$ is defined on some ray of the form $[T_y, +\infty)$ and for each $T \geq T_y$ it is fulfilled $\sup\{|y(t)| : t \geq T\} > 0$. Such a solution is called a *regular solution* of (E_1) .

We say that a real valued function u defined on an interval $[a, +\infty)$ has some property *eventually*, if there is a number $b \geq a$ such that u has this property on the interval $[b, +\infty)$.

A regular solution $y(t)$ of equation (E_1) is said to be *nonoscillatory*, if there exists a number $t_0 \geq 0$ such that $y(t)$ is of constant sign for every $t \geq t_0$. Otherwise, it is called *oscillatory*. Also, note that a *nonoscillatory* solution is called *eventually positive* (*eventually negative*), if the constant sign that determines

its *nonoscillation* is positive (negative). Equation (E_1) is called *oscillatory*, if all its solutions are oscillatory.

Moreover, in this article, when we write a functional relation (or inequality), we will mean that it holds for all sufficiently large values of the argument.

In order to assist our investigations on the oscillation of the equation (E_1) , we shall consider in the next section the delay impulsive differential equation with variable coefficients of the form

$$z'(t) + q(t)z(s(t)) = 0, \quad t \neq \tau_k \tag{E_2}$$

$$\Delta z(\tau_k) + q_{\tau_k}z(s(\tau_k)) = 0, \quad k \in N$$

and the corresponding to it inequalities

$$z'(t) + q(t)z(s(t)) \leq 0, \quad t \neq \tau_k \tag{N_{2,\leq}}$$

$$\Delta z(\tau_k) + q_{\tau_k}z(s(\tau_k)) \leq 0, \quad k \in N$$

and

$$z'(t) + q(t)z(s(t)) \geq 0, \quad t \neq \tau_k \tag{N_{2,\geq}}$$

$$\Delta z(\tau_k) + q_{\tau_k}z(s(\tau_k)) \geq 0, \quad k \in N,$$

under the hypotheses:

$$\mathbf{H}_2^* \quad s \in C^1(R^+, R^+), \quad s'(t) > 0, \quad \lim_{t \rightarrow +\infty} s(t) = +\infty, \quad \text{and } s(t) < t.$$

$$\mathbf{H}_3^* \quad q \in P_\tau C(R^+, R^+), \quad q_{\tau_k} \in R, \quad 1 > q_{\tau_k} \geq 0, \quad k \in N.$$

3. Some Useful Lemmas

Consider $y(t)$ as a solution of equation (E_1) and set the auxiliary function

$$z(t) = y(t) - c(t)y(h(t)), \quad \Delta z(\tau_k) = \Delta y(\tau_k) - c_{\tau_k} \Delta y(h(\tau_k)), \quad c_{\tau_k} = c(\tau_k), \quad k \in N. \tag{*}$$

We introduce two lemmas, which investigate the asymptotic behavior of the function $z(t)$, when $y(t)$ is a non-oscillatory solution of (E_1) . The first one is formulated and proved for an eventually positive solution $y(t)$ of the equation (E_1) .

Lemma 1. *Assume that the hypotheses $(H_1) - (H_5)$ are satisfied and $y(t)$ be an eventually positive solution of (E_1) . Then $z(t)$ is a decreasing eventually positive function of t with not strict down-jumps and $\lim_{t \rightarrow +\infty} z(t) = 0$ with*

$$\lim_{\tau_k \rightarrow +\infty} |\Delta z(\tau_k)| = 0.$$

Proof. Let $y(t)$ be an eventually positive solution of the equation (E_1) , i.e. $y(t)$ is a solution of (E_1) and there exists a number $T_0 > 0$ such that $y(t) > 0$ for $t \geq \rho(T_0)$. Then, from (E_1) and utilizing $(*)$ we have

$$z'(t) = -p(t)y(\sigma(t)), \quad t \neq \tau_k, \quad k \in N, \quad t \geq T_0, \tag{1}$$

$$\Delta z(\tau_k) = -p_{\tau_k}y(\sigma(\tau_k)), \quad k \in N, \quad \tau_k \geq T_0.$$

From (1), in view of (H_5) , it follows that $z(t)$ is an eventually decreasing function of t ($z'(t) < 0$) with not strict down-jumps ($\Delta z(\tau_k) \leq 0$) for $t \in [T_0, +\infty)$.

Assume $z(t) < 0$ eventually. Then, for some $t_1 \geq T_0$ there exists $\delta_\nu > 0$ such that $z(t) \leq -\delta_\nu$, for every $t \geq t_1, t \neq \tau_k$, i.e. $y(t) - c(t)y(h(t)) \leq -\delta_\nu, t \neq \tau_k, t \geq t_1$. In the meantime, for the same $\delta_\nu > 0$, there will be such a position ν in the sequence of the impulsive moments $\{\tau_k\}$, whereafter $z(\tau_k) \leq -\delta_\nu$, for every $\tau_k \geq \tau_\nu$ where $k \geq \nu, k \in N, \nu \in N$. Hence, $y(\tau_k) - c_{\tau_k}y(h(\tau_k)) \leq -\delta_\nu, \tau_k \geq \tau_\nu, k \geq \nu$. Denote $t_\nu = \max\{t_1, \tau_\nu\}$. Using (H_3) , we can combine the last two inequalities as

$$y(t) \leq -\delta_\nu + c(t)y(h(t)) \leq -\delta_\nu + c_0y(h(t)), \quad t \geq t_\nu.$$

By iterations, from the last inequality we get

$$y(t) \leq -\delta_\nu(1 + c_0 + c_0^2 + \dots + c_0^{n-1}) + c_0^n y(h^n(t)), \quad t \geq t_\nu, \tag{2}$$

In view of (H_3) , the inequality (2) implies

$$y(t) \leq -\frac{\delta_\nu}{1 - c}, \quad t \geq t_\nu.$$

This is a contradiction. Hence, our assumption, that eventually $z(t) < 0$, is impossible.

Assume $z(t) \equiv 0$. Then, from (1), it follows $p(t)y(\sigma(t)) \equiv 0$ and $p(\tau_k)y(\sigma(\tau_k)) = 0$. But $y(t)$ is an eventually positive function, so we should have $p(t) \equiv 0, p_{\tau_k} = 0, k \in N$, which contradicts (H_4) .

Thus, $z(t) \geq 0$ eventually. Moreover, in view of (1) and (H_5) , we conclude that there exists $\lim_{t \rightarrow +\infty} z(t)$ and it is a finite positive number or zero. Observe, that the last fact implies $\lim_{\tau_k \rightarrow +\infty} |\Delta z(\tau_k)| = 0$.

Assume $\lim_{t \rightarrow +\infty} z(t) = L, L > 0$. Then, if we integrate (E_1) from T_0 to t , we obtain

$$\int_{T_0}^t z'(s)ds + \int_{T_0}^t p(s)y(\sigma(s))ds = 0,$$

or

$$z(t) - z(T_0) - \sum_{T_0 < \tau_k < t} \Delta z(\tau_k) + \int_{T_0}^t p(s)y(\sigma(s))ds = 0,$$

i.e.

$$z(t) = z(T_0) + \sum_{T_0 < \tau_k < t} \Delta z(\tau_k) - \int_{T_0}^t p(s)y(\sigma(s))ds. \tag{3}$$

But $\Delta z(\tau_k) = -p_{\tau_k}y(\sigma(\tau_k))$ and from (3) we get

$$z(t) = z(T_0) - \sum_{T_0 < \tau_k < t} p_{\tau_k}y(\sigma(\tau_k)) - \int_{T_0}^t p(s)y(\sigma(s))ds. \tag{4}$$

Note, that $L < z(t) < y(t)$, i.e. $y(t)$ is bounded from below. Then (4) reduces to

$$z(t) \leq z(T_0) - L \left(\sum_{T_0 < \tau_k < t} p_{\tau_k} + \int_{T_0}^t p(s)ds \right),$$

which together with (H_5) implies $\lim_{t \rightarrow +\infty} z(t) = -\infty$ and contradicts our assumption. Therefore, $\lim_{t \rightarrow +\infty} z(t) = 0$. The proof is complete.

The second lemma is only formulated for an eventually negative solution $y(t)$ of the equation (E_1) , but the proof is carried out respectively to the proof of Lemma 1.

Lemma 2. *Assume that the hypotheses $(H_1) - (H_5)$ are satisfied. Let $y(t)$ be an eventually negative solution of (E_1) . Then $z(t)$ is an increasing eventually negative function of t with not strict up-jumps and $\lim_{t \rightarrow +\infty} z(t) = 0$ with $\lim_{\tau_k \rightarrow +\infty} |\Delta z(\tau_k)| = 0$.*

Our aim into the next lemma is to establish appropriate sufficient condition under which the equation (E_2) is oscillatory. To this end, we introduce the following result.

Lemma 3. *Assume the hypotheses $(H_1), (H_2^*), (H_3^*)$ are satisfied. Suppose also that:*

$$\frac{1}{e} \limsup_{t \rightarrow \infty} \prod_{s(t) < \tau_k < t} (1 - q_{\tau_k}) < \liminf_{t \rightarrow \infty} \int_{s(t)}^t q(r)dr, \quad k \in N.$$

Then:

- (a) the equation (E_2) is oscillatory;
- (b) the inequality $(N_{2,\leq})$ has no eventually positive solutions;
- (c) the inequality $(N_{2,\geq})$ has no eventually negative solutions.

Proof. Since the proofs of (a),(b) and (c) can be carried out by similar arguments, it suffices to prove only the case (a). To this end, we assume for the sake of contradiction, that equation (E_2) has a nonoscillatory solution. Since the negative of a solution of (E_2) is again a solution of (E_2) , it suffices to prove the lemma considering this solution as an eventually positive function. So, suppose that $z(t)$ is a solution of (E_2) and there exists a number $t_0 > 0$, such that $z(t) > 0$, for every $t \geq s(t_0)$. Then, it follows from (E_2) that $z'(t) = -q(t)z(s(t)) < 0$ and $\Delta z(\tau_k) = -q_{\tau_k}z(s(\tau_k)) \leq 0$, for every $t, \tau_k \geq s(t_0)$, $k \in N$, i.e. $z(t)$ is a decreasing function with not strict down-jumps.

Now, we can rearrange (E_2) , dividing by $z(t)$, in order to obtain

$$\frac{z'(t)}{z(t)} = -q(t)\frac{z(s(t))}{z(t)} < -q(t), \quad t \neq \tau_k, \quad k \in N, \tag{5}$$

$$\Delta z(\tau_k) = -q_{\tau_k}z(s(\tau_k)) < -q_{\tau_k}z(\tau_k), \quad k \in N.$$

It follows from the condition of the lemma, that there exist a constant $L > 0$ and a number $t_1 \geq t_0$, such that

$$\frac{\int_{s(t)}^t q(r)dr}{m} \geq L > \frac{1}{e}, \quad t \geq t_1, \tag{6}$$

where we denote

$$m = \limsup_{t \rightarrow \infty} \prod_{s(t) < \tau_k < t} (1 - q_{\tau_k}).$$

If we integrate (5) from $s(t)$ to t , we obtain

$$\int_{s(t)}^t \frac{z'(r)}{z(r)}dr < - \int_{s(t)}^t q(r)dr,$$

or

$$\ln \frac{z(t)}{z(s(t))} + \sum_{s(t) < \tau_k < t} \ln \frac{z(\tau_k)}{z(\tau_k + 0)} < - \int_{s(t)}^t q(r)dr. \tag{7}$$

Moreover, $z(\tau_k + 0) - z(\tau_k) = -q_{\tau_k}z(s(\tau_k)) < -q_{\tau_k}z(\tau_k)$ and $z(\tau_k + 0) < (1 - q_{\tau_k})z(\tau_k)$, i.e. $\frac{1}{1 - q_{\tau_k}} < \frac{z(\tau_k)}{z(\tau_k + 0)}$. So, $\ln \frac{1}{1 - q_{\tau_k}} < \ln \frac{z(\tau_k)}{z(\tau_k + 0)}$ and from (6) and (7) we get

$$\ln \left[\frac{z(t)}{z(s(t))} \prod_{s(t) < \tau_k < t} \frac{1}{1 - q_{\tau_k}} \right] < - \int_{s(t)}^t q(r)dr,$$

i.e.

$$\ln \left[\frac{z(s(t))}{z(t)} \prod_{s(t) < \tau_k < t} (1 - q_{\tau_k}) \right] > Lm.$$

Using the inequality $e^x > ex$, it follows from the last inequality that

$$\frac{z(s(t))}{z(t)} \prod_{s(t) < \tau_k < t} (1 - q_{\tau_k}) > eLm,$$

which implies

$$\frac{z(s(t))}{z(t)} > eL.$$

Repeating the above procedure by induction on (5), we conclude that there exists a sequence $\{t_n\}$ where $t_n \rightarrow \infty$ as $n \rightarrow \infty$, such that

$$\frac{z(s(t))}{z(t)} > (eL)^n, \quad t \geq t_n. \tag{8}$$

Choose n such that

$$\left(\frac{2}{mL} \right)^2 < (eL)^n, \tag{9}$$

which is possible because $eL > 1$, by (6). Further, fix arbitrary chosen \hat{t} , where $\hat{t} \geq t_n$.

It follows from (6), that there exists a number $\xi \in [s(\hat{t}), \hat{t}]$, such that

$$\int_{s(\hat{t})}^{\xi} q(r)dr \geq \frac{mL}{2}, \quad \int_{\xi}^{\hat{t}} q(r)dr > \frac{mL}{2}.$$

If we integrate (E_2) over the interval $[s(\hat{t}), \xi]$, we find

$$z(\xi) - z(s(\hat{t})) - \sum_{s(\hat{t}) < \tau_i < \xi} \Delta z(\tau_i) + \int_{s(\hat{t})}^{\xi} q(r)z(s(r))dr = 0,$$

or

$$z(\xi) - z(s(\hat{t})) + \sum_{s(\hat{t}) < \tau_i < \xi} q_{\tau_i} z(s(\tau_i)) + \int_{s(\hat{t})}^{\xi} q(s) z(s(r)) dr = 0.$$

By omitting the first and the third terms and using the decreasing nature of $z(t)$ we find

$$z(s(\hat{t})) > z(s(\xi)) \int_{s(\hat{t})}^{\xi} q(r) dr,$$

i.e.

$$\frac{z(s(\hat{t}))}{z(s(\xi))} > \frac{mL}{2}. \tag{10}$$

Similarly, integrating (E_2) over the interval $(\xi, \hat{t}]$, we find

$$z(\hat{t}) - z(\xi) - \sum_{\xi < \tau_i < \hat{t}} \Delta z(\tau_i) + \int_{\xi}^{\hat{t}} q(r) z(s(r)) dr = 0,$$

or

$$z(\hat{t}) - z(\xi) + \sum_{\xi < \tau_i < \hat{t}} q_{\tau_i} z(s(\tau_i)) + \int_{\xi}^{\hat{t}} q(r) z(s(r)) dr = 0.$$

By omitting the first and the third terms and using the decreasing nature of $z(t)$ we find

$$z(\xi) > z(s(\hat{t})) \int_{\xi}^{\hat{t}} q(r) dr,$$

i.e.

$$\frac{z(\xi)}{z(s(\hat{t}))} > \frac{mL}{2}. \tag{11}$$

From (10) and (11) we conclude

$$\frac{z(s(\xi))}{z(\xi)} < \left(\frac{2}{mL} \right)^2,$$

which, together with (8), imply

$$(eL)^n < \frac{z(s(\xi))}{z(\xi)} < \left(\frac{2}{mL} \right)^2. \tag{12}$$

Note that (12) is in contradiction with (9). The proof of the lemma is complete.

The last lemma in this section affords us new opportunity to investigate the oscillation of the equation (E_1) . It indicates, that the function $z(t)$, which is constructed from an eventually none-oscillatory solution $y(t)$ of equation (E_1) , is found to satisfy an auxiliary relation with useful properties. To this purpose, we introduce the following hypotheses, which describe the possible location of the points of impulse effect:

(AH) $\exists n \geq 1, n \in N : h(\tau_s) \in \{\tau_k\}_{k=1}^\infty, \tau_s \in \{\tau_i\}_{i=n+1}^\infty$.

(BH) There exists a strictly increasing sequence $\{k_\nu\}_{\nu=1}^\infty \subseteq N$ (not obligatory consistent), for which $\{\tau_{k_\nu}\}_{\nu=1}^\infty \subseteq \{\tau_k\}_{k=1}^\infty$, but $h(\tau_{k_\nu}) \notin \{\tau_k\}_{k=1}^\infty$.

Lemma 4. *Let the hypotheses $(H_1) - (H_5)$ are satisfied and $y(t)$ be a positive solution of the equation (E_1) . Then the function $z(t)$, defined by $(*)$, satisfies the neutral impulsive differential relation*

$$z'(t) - r(t)z'(h(t)) + p(t)z(\sigma(t)) = 0, t \neq \tau_k, k \in N \tag{E_1^*}$$

$$\Delta z(\tau_k) - r_{\tau_k} \Delta z(h(\tau_k)) + p_{\tau_k} z(\sigma(\tau_k)) \leq 0, k \in N,$$

where $r(t) = c(\sigma(t)) \frac{p(t)}{p(h(t))}, r_{\tau_k} = r(\tau_k) = c_{\sigma(\tau_k)} \frac{p_{\tau_k}}{p_{h(\tau_k)}}$.

Proof. A direct substitution shows that $z(t) = y(t) - c(t)y(t - h), t \neq \tau_k, k \in N$ is a continuously differentiable solution of the differential part of (E_1^*) in $J = \bigcup_{k=0}^{+\infty} (\tau_k, \tau_{k+1})$.

Using the definition of the function $z(t)$ in $(*)$, we can consider the left side of the difference part of (E_1^*) , as follows:

$$\begin{aligned} \Delta \{y(\tau_k) - c_{\tau_k} y(h(\tau_k))\} - r_k \Delta \{y(h(\tau_k)) - c_{h(\tau_k)} y(h^2(\tau_k))\} \\ + p_{\tau_k} \{y(\sigma(\tau_k)) - c_{\sigma(\tau_k)} y(\sigma(h(\tau_k)))\}. \end{aligned}$$

This, by means of the difference part of (E_1) , can be reduced to

$$-r_{\tau_k} \{ \Delta (y(h(\tau_k)) - c_{h(\tau_k)} y(h^2(\tau_k)) + p_{h(\tau_k)} y(\sigma(h(\tau_k)))) \} = I_k.$$

Observe, that when the hypothesis **(AH)** is valid, then $h(\tau_k)$ does belong to the sequence $\{\tau_k\}_{k=n+1}^\infty$. Therefore, we have eventually $I_k = 0$.

Assume, the hypothesis **(BH)** is valid. Then, $I_k = 0$ for every $k < k_1$, while when $k = k_1$, we have $\Delta y(h(\tau_{k_1})) = 0$ and subsequently, we obtain

$$I_k = I_{k_1} = r_{\tau_{k_1}} c_{h(\tau_{k_1})} \Delta y(h^2(\tau_{k_1})) - r_{\tau_{k_1}} p_{h(\tau_{k_1})} y(\sigma(h(\tau_{k_1}))).$$

But, in view of **(BH)**, $h^2(\tau_{k_1})$ is not a point of impulse effect, i.e. $h^2(\tau_{k_1})$ does not belong to the sequence $\{\tau_k\}_{k=1}^\infty$ and $\Delta y(h^2(\tau_{k_1})) = 0$. Indeed, let us suppose, that $h^2(\tau_{k_1}) \in \{\tau_k\}_{k=1}^\infty$, whereas $h(\tau_{k_1}) \notin \{\tau_k\}_{k=1}^\infty$. From the hypotheses (H_2) and **(BH)**, it follows $h^{-1}(\tau_i) \in \{\tau_k\}_{k=1}^{k_1}$, $i = 0, k_1 - 1$. Thus, we should have $h^{-1}(h^2(\tau_{k_1})) \in \{\tau_k\}_{k=1}^{k_1}$, i.e. $h(\tau_{k_1}) \in \{\tau_k\}_{k=1}^{k_1}$, which is a contradiction. Hence, for $k = k_1$ we have

$$I_k = -r_{\tau_k} p_{h(\tau_k)} y(\sigma(h(\tau_k))) \leq 0. \tag{13}$$

Further, we have $I_k = 0$, $k_1 < k < k_2$, whereas $\Delta y(h(\tau_{k_2})) = 0$ and subsequently

$$I_k = I_{k_2} = r_{\tau_{k_2}} c_{h(\tau_{k_2})} \Delta y(h^2(\tau_{k_2})) - r_{\tau_{k_2}} p_{h(\tau_{k_2})} y(\sigma(h(\tau_{k_2}))).$$

Here, analogically as above, we see that $h^2(\tau_{k_2})$ is not a point of impulse effect, i.e. $h^2(\tau_{k_2})$ does not belong to the sequence $\{\tau_k\}_{k=1}^\infty$ i.e. $\Delta y(h^2(\tau_{k_2})) = 0$ and (13) is fulfilled, for $k = k_2$. This procedure can be repeated for every $k \in \{k_\nu\}_{\nu=1}^\infty \subseteq N$.

So, we conclude that under the conditions of the lemma, the difference part of (E_1^*) is satisfied by $\Delta z(\tau_k) = \Delta y(\tau_k) - c_{\tau_k} \Delta y(h(\tau_k))$, as well. The proof of the lemma is complete.

4. Oscillation Criteria for the Solutions of (E_1)

In this section we study the oscillatory properties of the equation (E_1) . The next theorems will establish sufficient conditions for oscillation of (E_1) .

Theorem 1. *Let the hypotheses $(H_1) - (H_5)$ are satisfied, $p_{\tau_k} \in (0, 1)$, $k \in N$ and*

$$(i) \quad \liminf_{k \rightarrow \infty} \frac{\sum_{\tau_k < \tau_i < \sigma^{-1}(\tau_k)} p_i + \int_{\tau_k}^{\sigma^{-1}(\tau_k)} p(s) ds}{1 - p_{\tau_k}} \geq 1.$$

Then, all solutions of equation (E_1) are oscillatory.

Proof. Assume, for the sake of contradiction, that (E_1) has a non-oscillatory solution. Since the negative of a solution of (E_1) is again a solution of (E_1) , it suffices to prove the theorem considering an eventually positive solution of (E_1) . So, let us suppose that $y(t)$ is a solution of (E_1) and there exists a number $T_0 > 0$, such that $y(t) > 0$, for $t \geq \rho(T_0)$. Recall $(*)$. By Lemma 4, $z(t)$

satisfies (E_1^*) . Moreover, from Lemma 1, it follows that $z(t)$ is an eventually non-increasing positive function of t with down-jumps, i.e.

$$z(t) > 0, \quad z'(t) \leq 0, \quad t \geq T_0, \quad \Delta z(\tau_k) \leq 0, \quad \tau_k \geq T_0, \quad k \in N. \quad (19)$$

In view of (19), the relation (E_1^*) leads us to the conclusion, that $z(t)$ satisfies the impulsive differential inequality with retarded argument of the form

$$z'(t) + p(t)z(\sigma(t)) \leq 0, \quad t \neq \tau_k, \quad t \geq T_0, \quad (20)$$

$$\Delta z(\tau_k) + p_{\tau_k}z(\sigma(\tau_k)) \leq 0, \quad \tau_k \geq T_0, \quad k \in N.$$

In view of (H_2) , we can integrate (20) from τ_k to $\sigma^{-1}(\tau_k)$, in order to find

$$z(\sigma^{-1}(\tau_k)) - z(\tau_k + 0) - \sum_{\tau_k < \tau_i < \sigma^{-1}(\tau_k)} \Delta z(\tau_i) + \int_{\tau_k}^{\sigma^{-1}(\tau_k)} p(s)z(\sigma(s))ds \leq 0,$$

i.e.

$$z(\sigma^{-1}(\tau_k)) - z(\tau_k + 0) + \sum_{\tau_k < \tau_i < \sigma^{-1}(\tau_k)} p_{\tau_i}z(\sigma(\tau_i)) + \int_{\tau_k}^{\sigma^{-1}(\tau_k)} p(s)z(\sigma(s))ds \leq 0.$$

Since $z(t)$ is an eventually decreasing function, by (19), it follows from the last inequality

$$z(\sigma^{-1}(\tau_k)) - z(\tau_k + 0) + z(\tau_k) \sum_{\tau_k < \tau_i < \sigma^{-1}(\tau_k)} p_{\tau_i} + z(\tau_k) \int_{\tau_k}^{\sigma^{-1}(\tau_k)} p(s)ds \leq 0. \quad (21)$$

Moreover, we have $\Delta z(\tau_k) \leq -p_{\tau_k}z(\sigma(\tau_k)) \leq -p_{\tau_k}z(\tau_k)$, $\tau_k \geq T_0$, $k \in N$.

That is why $z(\tau_k + 0) \leq (1 - p_{\tau_k})z(\tau_k)$, i.e. $\frac{z(\tau_k + 0)}{1 - p_{\tau_k}} \leq z(\tau_k)$. Thus,

(21) yields

$$z(\sigma^{-1}(\tau_k)) + z(\tau_k + 0) \left(\frac{\sum_{\tau_k < \tau_i < \sigma^{-1}(\tau_k)} p_i + \int_{\tau_k}^{\sigma^{-1}(\tau_k)} p(s)ds}{1 - p_{\tau_k}} - 1 \right) \leq 0.$$

Since $z(t)$ is an eventually positive function, the last inequality implies

$$\frac{\sum_{\tau_k < \tau_i < \sigma^{-1}(\tau_k)} p_i + \int_{\tau_k}^{\sigma^{-1}(\tau_k)} p(s)ds}{1 - p_{\tau_k}} < 1.$$

This contradicts the condition (i) of the theorem. Therefore, it is impossible for (20) to have an eventually positive solution. This contradicts (19) and completes the proof.

Theorem 2. *Let the hypotheses $(H_1) - (H_5)$ are satisfied. Suppose also that:*

(iv) $\sigma(t) \leq h(t) < t, \quad t \in R^+;$
 (v) $\exists r_0 > 0, : r(t) = c(\sigma(t)) \frac{p(t)}{p(h(t))} \geq r_0, r_{\tau_k} = c_{\sigma(\tau_k)} \frac{p_{\tau_k}}{p_{h(\tau_k)}} \geq r_0, k \in N;$

(vi) $\liminf_{t \rightarrow +\infty} [\int_t^{h^{-1}(t)} p(r)dr + \sum_{t < \tau_k < h^{-1}(t)} p_{\tau_k}] \geq 1 + r_0.$

Then, the equation (E_1) is oscillatory.

Proof. Assume for the sake of contradiction, that $y(t)$ is an eventually positive solution of the equation (E_1) . Then, in view of Lemma 1 and Lemma 4, the function $z(t)$, defined by $(*)$, is a decreasing eventually positive function with down-jumps that satisfies (E_1^*) . That is, $z(t)$ satisfies

$$z'(t) - r(t)z'(h(t)) + p(t)z(\sigma(t)) = 0, \quad t \neq \tau_k \tag{22}$$

$$\Delta z(\tau_k) - r_{\tau_k} \Delta z(h(\tau_k)) + p_{\tau_k} z(\sigma(\tau_k)) \leq 0, \quad k \in N.$$

In view of (H_2) , we can integrate (22) from t to $h^{-1}(t)$. Thus, we have

$$z(h^{-1}(t)) - z(t) - \sum_{t < \tau_k < h^{-1}(t)} \Delta z(\tau_k) - \int_t^{h^{-1}(t)} r(s)z'(h(s))ds + \int_t^{h^{-1}(t)} p(s)z(\sigma(s))ds = 0.$$

Now, using the condition (v) of the theorem, we obtain

$$z(h^{-1}(t)) - z(t) - r_0 z(t) + r_0 z(h(t)) - \sum_{t < \tau_k < h^{-1}(t)} [\Delta z(\tau_k) - r_0 \Delta z(h(\tau_k))]$$

$$+ \int_t^{h^{-1}(t)} p(s)z(\sigma(s))ds \leq 0.$$

By omitting the first and the fourth terms in the last inequality, we derive

$$-z(t) - r_0z(t) + \sum_{t < \tau_k < h^{-1}(t)} p_{\tau_k}z(\sigma(\tau_k)) + \liminf_{t \leq s \leq h^{-1}(t)} z(\sigma(s)) \int_t^{h^{-1}(t)} p(s)ds < 0, \tag{23}$$

because

$$\Delta z(\tau_k) - r_0\Delta z(h(\tau_k)) \leq \Delta z(\tau_k) - r_k\Delta z(h(\tau_k)) \leq -p_{\tau_k}z(\sigma(\tau_k)), \quad k \in N.$$

From Lemma 1 and the condition (iv) of the theorem in view of (H₂), it follows

$$\liminf_{t \leq s \leq h^{-1}(t)} z(\sigma(s)) \geq z(t) \quad \text{and} \quad z(\sigma(\tau_k)) > z(t) \quad \text{for} \quad t < \tau_k < h^{-1}(t). \tag{24}$$

Finally, counting on (24), we get from (23)

$$z(t) \left(-1 - r_0 + \sum_{t < \tau_k < h^{-1}(t)} p_{\tau_k} + \int_t^{h^{-1}(t)} p(r)dr \right) < 0,$$

which implies

$$\int_t^{h^{-1}(t)} p(r)dr + \sum_{t < \tau_k < h^{-1}(t)} p_{\tau_k} < 1 + r_0. \tag{25}$$

The conclusion obtained in (25) contradicts to the condition (vi) of the theorem. The proof is complete.

Theorem 3. *Let the hypotheses (H₁) – (H₅) are satisfied. Suppose also that:*

- (vii) $\sigma(t) < h(t) < t, \quad t \in R^+;$
- (viii) $\exists r_0 < 1 : r(t) = c(\sigma(t)) \frac{p(t)}{p(h(t))} \leq r_0 \quad \text{and} \quad r'(t) \geq 0, \quad t \in R^+;$
- (ix) $p_{\tau_k} < 1, \quad k \in N ;$
- (x) $\frac{1}{e} \limsup_{t \rightarrow \infty} \left(\prod_{h^{-1}(\sigma(t)) < \tau_k < t} (1 - p_{\tau_k}) \right) < \liminf_{t \rightarrow \infty} \int_{h^{-1}(\sigma(t))}^t \frac{p(s)}{1 - r(h^{-1}(\sigma(s)))} ds, \quad k \in N.$

Then, the equation (E_1) is oscillatory.

Proof. Assume for the sake of contradiction, that $y(t)$ is an eventually positive solution of the equation (E_1) . Then, in view of Lemma 1, the function $z(t)$, defined by $(*)$, is a decreasing to zero eventually positive function with down-jumps, i.e.

$$z(t) > 0, \quad \Delta z(\tau_k) \leq 0, \quad z'(t) < 0, \quad \lim_{t \rightarrow +\infty} z(t) = 0 \quad (26)$$

Let set a second auxiliary function

$$w(t) = z(t) - r(t)z(h(t)), \quad \Delta w(\tau_k) = \Delta z(\tau_k) - r_{\tau_k} \Delta z(h(\tau_k)), \quad k \in N, \quad (**)$$

where $r(t) = c(\sigma(t)) \frac{p(t)}{p(h(t))}$, $r_{\tau_k} = c_{\sigma(\tau_k)} \frac{p_{\tau_k}}{p_{h(\tau_k)}}$.

Obviously, we have

$$w'(t) = z'(t) - r(t)z'(h(t)) - r'(t)z(h(t)). \quad (27)$$

But, $z(t)$ satisfies (E_1^*) , by Lemma 4. Therefore, (27) yields

$$w'(t) = -p(t)z(\sigma(t)) - r'(t)z(h(t)) < 0.$$

Moreover, $\Delta w(\tau_k) = \Delta z(\tau_k) - r_{\tau_k} \Delta z(h(\tau_k)) = -p_{\tau_k} z(h(\tau_k)) < 0$. Hence, $w(t)$ is a decreasing function with down-jumps. This means that $L = \lim_{t \rightarrow +\infty} w(t)$ does exist.

Assume $L < 0$, i.e. $w(t)$ is an eventually negative function. Then, by condition $(\ddot{v}iii)$, we have $0 > w(t) = z(t) - r(t)z(h(t)) \geq z(t) - r_0 z(h(t))$, which in view of (26) and the monotonicity of $w(t)$, leads us to a contradiction. Therefore, $L \geq 0$ and we conclude, that

$$w(t) \geq 0, \quad w(t) \leq z(t). \quad (28)$$

Further, using the fact, that $1 - r(t) > 0$, utilizing the Lemma 1 and counting on the decreasing positive nature of $w(t)$, we find $w(t) = z(t) - r(t)z(h(t)) \leq (1 - r(t))z(h(t))$, which can be rewritten as

$$w(h^{-1}(\sigma(t))) \leq (1 - r(h^{-1}(\sigma(t))))z(\sigma(t)).$$

If we multiply the last inequality by $\frac{-p(t)}{1 - r(h^{-1}(\sigma(t)))} < 0$, we obtain

$$\frac{-p(t)}{1 - r(h^{-1}(\sigma(t)))} w(h^{-1}(\sigma(t))) \geq -p(t)z(\sigma(t)) = w'(t) + r'(t)z(h(t)),$$

or more precisely

$$w'(t) + \frac{p(t)}{1 - r(h^{-1}(\sigma(t)))} w(h^{-1}(\sigma(t))) \leq -r'(t)z(h(t)) < 0. \quad (29)$$

Moreover, from (E_1^*) , (28), the condition (vii) and the monotonicity of $w(t)$, it follows

$$\Delta w(\tau_k) = -p_{\tau_k} z(\sigma(\tau_k)) \leq -p_{\tau_k} w(\sigma(\tau_k)) < -p_{\tau_k} w(h^{-1}(\sigma(\tau_k))), \quad k \in N. \quad (30)$$

Finally, from (29) and (30) we can conclude, that the eventually positive function $w(t)$ satisfies the delay impulsive differential inequality

$$w'(t) + \frac{p(t)}{1 - r(h^{-1}(\sigma(t)))} w(h^{-1}(\sigma(t))) < 0$$

$$\Delta w(\tau_k) + p_{\tau_k} w(h^{-1}(\sigma(\tau_k))) < 0, \quad k \in N,$$

which has the form of $(N_{2,\leq})$. But, the conclusion obtained, under the conditions of the theorem and in view of Lemma 3(b), contradicts to (28). The proof is complete.

Corollary 1. *Let the conditions of Theorem 1, or Theorem 2, or Theorem 3 hold. Then:*

- (i) *the inequality $(N_{1,\leq})$ has no eventually positive solutions;*
- (ii) *the inequality $(N_{1,\geq})$ has no eventually negative solutions.*

The proof of the corollary is carried out analogously to the proofs of the respective theorems.

References

- [1] D.D. Bainov, M.B. Dimitrova, Sufficient conditions for oscillation of the solutions of a class of impulsive differential equations with deviating arguments, *J. Appl. Math., Stochastic Analysis*, **9** (1996), 33-42.
- [2] M.B. Dimitrova, Criteria for oscillation of impulsive differential equations of first order with deviating arguments, *Kyungpook Math. J.*, **40** (2000), 29-37.
- [3] M.B. Dimitrova, V.I. Donev, Existence and asymptotic behavior of the positive solutions of neutral impulsive differential equations, *Nonlinear Oscillations*, Kiev, **8**, No. 3 (2005), 304-318.

- [4] M.B. Dimitrova, V.I. Donev, Oscillatory properties of first order neutral delay impulsive differential equations with constant coefficients, In: *Proceedings of 1-st International Conference AMiTaNS'09* (Ed-s: M.D. Todorov, C.I. Christov), American Institute of Physics CP1186 (2009), 259-270.
- [5] A. Dishliev, K. Dishlieva, S. Nenov, *Specific Asymptotic Properties of the Solutions of Impulsive Differential Equations. Methods and Applications*, Academic Publications, Ltd (2011).
- [6] L.H. Erbe, Q. Kong, B.G. Zhang, *Oscillation Theory for Functional Differential Equations*, Marcel Decker Inc., New York (1995).
- [7] J.R. Graef, J.H. Shen, I.P. Stavroulakis, Oscillation of impulsive neutral delay differential equations, *J. Math. Anal. Appl.*, **268**, No. 1 (2002), 310-333.
- [8] M.K. Grammatikopoulos, M.B. Dimitrova, V.I. Donev, Oscillations of first order delay impulsive differential equations, *Technical Report* (Ed-s: I.P. Stavroulakis at al), University of Ioannina, Greece, **16** (2007), 171-182.
- [9] M.K. Grammatikopoulos, M.B. Dimitrova, V.I. Donev, *Oscillations of first order impulsive differential equations with variable coefficients and advanced argument*, In: *Proceedings of 33-th International Conference AMEE'33* (Ed. M.D. Todorov), American Institute of Physics, CP946 (2007), 206-214.
- [10] K. Gopalsamy, B.G. Zhang, On delay differential equations with impulses, *J. Math. Anal. Appl.*, **139** (1989), 110-122.
- [11] M.K. Grammatikopoulos, E.A. Grove, G. Ladas, Oscillation and asymptotic behavior of neutral differential equations with deviating arguments, *Applicable Analysis*, **22** (1986), 1-19.
- [12] I. Gyori, G. Ladas, *Oscillation Theory of Delay Differential Equations with Applications*, Clarendon Press, Oxford (1991).
- [13] A.D. Myshkis, *Linear Differential Equations with Retarded Argument*, Moskow, Nauka (1972), In Russian.
- [14] Y. Zhang, A. Zhao, J. Yan, Oscillation criteria for impulsive delay differential equations, *J. Math. Anal. Appl.*, **205** (1997), 461-470.

- [15] A. Zhao, J. Yan, Existence of positive solutions of delay differential equations with impulses, *J. Math. Anal. Appl.*, **210** (1997), 667-678.