

**ON THE GEOMETRY OF SMALL WEIGHT CODEWORDS
OF DUAL ALGEBRAIC GEOMETRIC CODES**

Claudio Fontanari¹ §, Chiara Marcolla²

Dipartimento di Matematica

Università di Trento

Via Sommarive 14, 38123 Trento, ITALY

Abstract: We investigate the geometry of the support of small weight codewords of dual algebraic geometric codes on smooth complete intersections by applying the powerful tools recently developed by Alain Couvreur. In particular, by restricting ourselves to the case of Hermitian codes, we recover and extend previous results obtained by the second named author joint with Marco Pellegrini and Massimiliano Sala.

AMS Subject Classification: 68Pxx

Key Words: small weight codewords, algebraic geometric codes

1. Introduction

In the recent contribution [3], the number of small weight codewords for some families of Hermitian codes is determined. Besides explicit computation, the main ingredient in [3] is a nice geometric characterization of the points in the support of a minimum weight codeword, which turn out to be collinear (see Corollary 1 and Proposition 2 in [3]).

Here we show that such a property is not peculiar to Hermitian codes, but it holds in full generality for dual algebraic geometric codes on any smooth

Received: April 17, 2012

© 2015 Academic Publications, Ltd.
url: www.acadpubl.eu

§Correspondence author

complete intersection projective variety of arbitrary dimension. Namely, by exploiting the algebro-geometric tools provided by [1], we prove the following:

Theorem 1. *Let $X \subset \mathbb{P}^r$, $r \geq 2$, be a smooth connected complete intersection defined over \mathbb{F}_q . Let G be a divisor on X such that $L(G) \supseteq H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(m))$. Let P_1, \dots, P_n be distinct rational points on X avoiding the support of G and let $D := P_1 + \dots + P_n$. Let d be the minimum distance of the code $C(D, G)$ and let $\{P_{i_1}, \dots, P_{i_d}\}$ be the points in the support of a minimum weight codeword.*

- (i) *If $d \leq m + 2$, then $d = m + 2$ and all the $m + 2$ points P_{i_j} are collinear in \mathbb{P}^r .*
- (ii) *If $d \leq 2m + 2$ and no $m + 2$ of the P_{i_j} 's are collinear, then $d = 2m + 2$ and all the $2m + 2$ points P_{i_j} lie on a plane conic.*
- (iii) *If $d \leq 3m$, no $m + 2$ of the P_{i_j} 's are collinear and no $2m + 2$ of them lie on a plane conic, then $d = 3m$ and all the $3m$ points P_{i_j} lie at the intersection of two complanar plane curves of respective degrees 3 and m .*

If we focus on the explicit interesting example of Hermitian codes (as presented for instance in [2]), then our general result specializes as follows:

Corollary 1. *Let $X \subset \mathbb{P}^2$ be the Hermitian curve of affine equation $x^{q+1} = y^q + y$ defined over \mathbb{F}_{q^2} . Let $G = \rho P_0$, where P_0 is the point at infinity, $D = P_1 + \dots + P_n$ with $n = q^3$, and let $C(D, G)$ be the Hermitian code. Let d be the minimum distance and let $\{P_{i_1}, \dots, P_{i_d}\}$ be the points in the support of a minimum weight codeword.*

- *If $0 \leq \rho \leq q^2 - q - 2$, then all the points P_{i_j} are collinear.*
- *If $\rho \geq q^2 - q - 2$, then the following holds.*
 - (i) *If $d \leq q$, then all the points P_{i_j} are collinear.*
 - (ii) *If $d \leq 2q - 2$, then either q of the points P_{i_j} are collinear, or all the points P_{i_j} lie on a plane conic.*
 - (iii) *If $d \leq 3q - 6$, then either q of the points P_{i_j} are collinear, or $2q - 2$ of the points P_{i_j} lie on a plane conic, or all the points P_{i_j} lie at the intersection of two complanar plane curves of respective degrees 3 and $q - 2$.*
- *If $\rho \geq q^2 - 1$, then the following holds.*
 - (i) *If $d \leq q + 1$, then all the points P_{i_j} are collinear.*

- (ii) If $d \leq 2q$, then either $q + 1$ of the points P_{i_j} are collinear, or all the points P_{i_j} lie on a plane conic.
- (iii) If $d \leq 3q - 3$, then either $q + 1$ of the points P_{i_j} are collinear, or $2q$ of the points P_{i_j} lie on a plane conic, or all the points P_{i_j} lie at the intersection of two coplanar plane curves of respective degrees 3 and $q - 1$.

In the case of Hermitian codes we may even describe the geometry of small, even not minimum, weight codewords (notice that our technical assumption $L(G) = H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d - 2))$ is satisfied by the so-called corner codes according to the terminology of [3], Definition 2):

Proposition 1. *Let $X \subset \mathbb{P}^2$ be the Hermitian curve of affine equation $x^{q+1} = y^q + y$ defined over \mathbb{F}_{q^2} . Let $G = \rho P_0$, where P_0 is the point at infinity, $D = P_1 + \dots + P_n$ with $n = q^3$, and let $C(D, G)$ be the Hermitian code. If $0 \leq \rho \leq q^2 - q - 2$ and $L(G) = H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d - 2))$, then at least $d - 1$ of the points $\{P_{i_1}, \dots, P_{i_{d+a}}\}$ in the support of a codeword of weight $d + a$, where $0 \leq a \leq d - 3$, are collinear.*

The present research is part of the PhD program of the second named author, supervised by Massimiliano Sala. The first named author has been partially supported by GNSAGA of INdAM and MIUR Cofin 2008 - Geometria delle varietà algebriche e dei loro spazi di moduli (Italy).

2. The Proofs

Proof of Theorem 1. Let $c \in C(D, G)$ be a minimum weight codeword having support $\{P_{i_1}, \dots, P_{i_s}\}$ with $s = d$. By the definition of a dual code, we have

$$\sum_{j=1}^s c_j f(P_{i_j}) = 0$$

for every $f \in L(G) \supseteq H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(m))$. In particular, we have

$$\sum_{j=1}^s c_j \text{ev}_{P_{i_j}}(f) = 0$$

for every $f \in H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(m))$. Hence $\text{ev}_{P_{i_j}}$ turn out to be linearly dependent in $H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(m))$ and, in the terminology of [1], Definition 2.4 and Definition

2.8, P_{i_1}, \dots, P_{i_s} are m -linked. Now, we have $d = s = m + 2$ in case (i) by [1], Proposition 6.1, $d = s = 2m + 2$ in case (ii) by [1], Proposition 7.2, and $d = s = 3m$ in case (iii) by [1], Proposition 8.1. Finally, we complete case (i) by [1], Proposition 6.2, case (ii) by [1], Proposition 7.4, and case (iii) by [1], Proposition 9.1.

□

Proof of Corollary 1. Since $L(G) = \langle \{x^i y^j : i \geq 0, 0 \leq j \leq q - 1, iq + j(q + 1) \leq \rho\} \rangle$ and $H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(k))$ can be identified with the set of polynomials in r variables of degree at most k , it is easy to check that $L(G) \supseteq H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(k))$ for every $k \geq 0$ such that $k \leq q - 1$ and $k(q + 1) \leq \rho$.

If $0 \leq \rho \leq q^2 - q - 2$, then let $m := d - 2$. By [2], §5.3, we have $\rho = 2q^2 - q - uq - v - 1$ with $1 \leq u, v \leq q - 1$ and $d = (q - u)q - v$ if $u < v$, $d = (q - u)q$ if $u \geq v$. Hence $d = m + 2$ implies $m = u - 1$ for $u > v$ and $m = u$ for $u = v$, hence

$$m(q + 1) \leq \rho. \tag{1}$$

If $\rho \geq q^2 - q - 2$ let $m := q - 2$, if instead $\rho \geq q^2 - 1$ let $m := q - 1$. In both cases (1) is satisfied, implying $L(G) \supseteq H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(m))$. Now our claim follows from Theorem 1.

□

Proof of Proposition 1. If c_j are the non-zero components of the corresponding codeword, then

$$\sum_{j=1}^{d+a} c_j f(P_{i_j}) = 0$$

for every $f \in L(G) \supseteq H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d - 2))$, in particular $\{P_{i_1}, \dots, P_{i_{d+a}}\}$ are $(d - 2)$ -linked.

If they are not minimally $(d - 2)$ -linked, then (up to reordering) we have

$$\sum_{j=1}^{d+a-1} b_j f(P_{i_j}) = 0$$

for every $f \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d - 2))$. Our assumption $L(G) = H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d - 2))$ implies that the b_j 's are the components of a codeword of weight strictly less than $d + a$. By induction on a starting from Corollary 1 we conclude that at least $d - 1$ of them are collinear.

Assume now that the points $\{P_{i_1}, \dots, P_{i_{d+a}}\}$ are minimally $(d - 2)$ -linked. If they are not collinear, there exists a hyperplane H containing exactly l of them, with $2 \leq l \leq d + a - 1$. By [1], Lemma 5.1, the remaining $d + a - l$ points are (minimally) $(d - 3)$ -linked and from [1], Proposition 7.2, it follows that at

least $(d - 1)$ of them are collinear since by our numerical assumption on a we have $d + a - l \leq 2(d - 3) + 1$. \square

References

- [1] A. Couvreur, The dual minimum distance of arbitrary dimensional algebraic-geometric codes, *J. Algebra*, **350** (2012), 84-107.
- [2] T. Høholdt, J. H. van Lint, and R. Pellikaan, *Algebraic Geometry of Codes*, Handbook of Coding Theory (1998), 871-961.
- [3] M. Pellegrini, C. Marcolla, M. Sala, On the weight of affine-variety codes and some Hermitian codes, In: *Proceedings of WCC 2011* (2011), 273-282.

